Table 4A

2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years (2)

Page: 1
Run Date: 7 / 8 / 04

Comparison of Racial and Ethnic Distribution by Clinic

Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other / Unknown %
(Delmar) P	1,556	3.9	90.9	2.0	0.2	1.3	1.7
(Maplewood	569	43.9	47.6	2.5	0.0	3.0	3.0
00Family C	1	*	*	*	*	*	*
00Miller C	127	95.3	0.0	3.1	0.0	0.0	1.6
01SL Conne	455	0.9	95.4	2.6	0.0	0.0	1.1
02Family C	552	12.9	69.0	1.8	0.2	2.5	13.6
Andrew Co	354	96.3	0.3	1.4	0.0	0.3	1.7
Atchison C	178	94.4	2.2	2.8	0.0	0.6	0.0
Audrain Co	576	78.3	12.7	5.0	0.2	0.2	3.6
Barnes Hos	168	12.5	61.9	19.0	0.0	3.0	3.6
Barry Co H	1,500	71.0	0.6	25.5	0.0	0.6	2.3
Barton Co	480	92.9	0.0	4.2	0.4	0.0	2.5
Bates Co H	525	83.2	1.5	4.2	0.6	0.0	10.5
BeltonCass	115	73.9	4.3	20.0	0.0	0.0	1.7
Benton Co	429	92.8	0.0	4.2	0.5	0.2	2.3
Blue Sprin	1,335	81.1	6.3	6.5	0.7	0.7	4.6
Bollinger	391	95.7	0.5	1.3	0.3	0.8	1.5
BransTaney	1,080	85.3	0.8	10.3	0.4	0.6	2.7
Butler Co	1,645	87.7	9.4	1.9	0.1	0.1	0.7

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with errors.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

CDC

2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years (2)

	Comparison of Racial and Ethnic Distribution by Clinic					
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other / Unknown %
Callaway C	651	82.0	11.2	4.1	0.3	0.5	1.8
Camden Co	896	90.4	0.2	6.1	0.2	0.4	2.6
Cape Girar	1,361	72.4	19.8	6.0	0.3	0.3	1.3
Carondelet	1,244	57.3	24.7	7.6	0.1	2.3	8.0
Carroll Co	308	95.1	2.3	1.6	0.0	0.0	1.0
Carter Co	265	97.4	0.0	0.8	0.0	0.0	1.9
Cass Co He	1,164	90.5	1.8	3.4	0.7	0.4	3.2
Cedar Co H	482	91.5	0.2	1.7	0.8	0.2	5.6
Chariton C	151	88.7	7.3	2.6	0.0	0.0	1.3
Christian	1,442	94.7	0.5	2.5	0.6	0.1	1.7
Clark Co H	207	95.7	0.5	1.9	0.0	0.0	1.9
Clay Chd-E	61	*	*	*	*	*	*
Clay Chd-P	50	*	*	*	*	*	*
Clay Co He	2,344	78.8	6.3	8.5	0.9	2.0	3.4
Clay Metro	17	*	*	*	*	*	*
Clinton Co	573	85.9	2.8	4.7	0.7	0.7	5.2
Cole Co He	1,781	68.6	21.3	6.7	0.2	0.4	2.8
Columbia-B	2,055	56.2	26.2	7.3	0.5	3.2	6.7
Comp Healt	735	0.7	96.5	1.4	0.0	0.0	1.5

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with errors.
 *   Percentages are not calculated if < 100 records are available for analysis after exclusions.

CDC

2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Racial and Ethnic Distribution by Clinic						
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other / Unknown %
Cooper Co	339	80.2	9.1	4.1	0.0	0.3	6.2
Crawford C	718	92.5	0.3	6.0	0.6	0.0	0.7
Crescent C	1,503	28.3	52.3	10.8	0.5	1.8	6.3
Dade Co He	276	92.8	1.4	3.3	0.0	0.4	2.2
Dallas Co	410	95.4	0.0	3.9	0.0	0.0	0.7
Daviess Co	284	93.3	0.4	3.2	0.0	0.7	2.5
Dekalb Co-	166	91.0	1.8	1.2	0.0	0.0	6.0
Dent Co He	521	95.8	0.6	1.9	0.6	0.0	1.2
Douglas Co	418	98.1	0.0	1.0	0.5	0.0	0.5
Dunklin Co	1,082	66.3	12.8	19.7	0.0	0.4	0.8
Elsber-Lnc	51	*	*	*	*	*	*
Florence H	891	0.8	97.4	1.2	0.0	0.0	0.6
Forest Par	751	16.2	76.6	1.7	0.0	0.3	5.2
Franklin C	1,597	93.7	2.1	2.3	0.1	0.6	1.3
FtWood-Pul	1,064	58.6	20.7	13.2	1.3	2.5	3.8
Gasconade	366	95.6	0.5	1.9	0.0	0.0	1.9
GascondOsg	161	92.5	0.6	2.5	0.0	0.0	4.3
Grace Hill	340	6.5	87.4	3.2	0.0	0.3	2.6
Grundy Co	362	90.6	0.8	7.5	0.0	0.0	1.1

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with errors.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 4A
2003 Pediatric Nutrition Surveillance [1]
Missouri
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years [2]

Comparison of Racial and Ethnic Distribution by Clinic						
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other / Unknown %
Hamilton/C	111	94.6	0.9	2.7	0.0	0.0	1.8
Harrison C	297	96.0	0.7	2.7	0.0	0.0	0.7
Hartville-	152	95.4	0.0	2.0	0.7	0.7	1.3
Henry Co H	647	91.5	2.6	1.4	0.0	0.5	4.0
HGP-Connec	178	0.0	96.6	1.7	0.0	0.0	1.7
Hickory Co	193	95.9	0.5	1.6	0.5	0.5	1.0
Holt Count	103	96.1	0.0	1.9	0.0	1.0	1.0
Hosphill-T	2,767	15.7	65.8	13.9	0.2	1.4	3.1
Howard Co	175	69.1	12.0	2.3	0.0	0.0	16.6
Iron Co He	405	95.1	3.2	1.5	0.0	0.0	0.2
Jfcac-Arno	1,361	94.3	1.5	1.7	0.1	0.3	2.1
Jfcac-Unit	1,706	94.5	2.3	1.7	0.2	0.6	0.5
John C Mur	2,540	16.5	70.9	7.3	0.1	1.8	3.4
Johnson Co	1,423	77.2	10.3	7.4	0.1	1.3	3.8
Joplin Cit	4,015	78.3	2.8	15.6	0.7	0.3	2.3
Kansas Cit	2	*	*	*	*	*	*
Kchd-Ruski	2	*	*	*	*	*	*
Knox Co He	151	94.7	0.0	4.0	0.0	0.7	0.7
La Clinica	849	30.0	28.7	8.8	0.6	8.5	23.3

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with errors.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 4A
2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Racial and Ethnic Distribution by Clinic						
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other / Unknown %
Laclede Co	1,017	92.5	0.8	3.3	0.2	0.5	2.7
Lakewood-T	1,638	70.9	9.3	13.2	0.7	1.2	4.8
Lawrence C	1,232	92.0	0.2	6.5	0.2	0.1	1.0
Lewis Co H	315	92.1	4.8	1.9	0.0	0.0	1.3
Lincoln Co	782	88.7	3.2	5.1	0.1	0.4	2.4
Linn Co He	360	89.7	1.7	3.9	0.3	0.3	4.2
Livingston	428	91.6	3.7	2.6	0.2	0.0	1.9
Macon Co H	408	82.1	7.8	1.7	0.2	0.2	7.8
Madison Co	546	92.5	0.5	4.6	0.4	0.2	1.8
Malden-Dnk	548	79.4	17.2	3.3	0.0	0.0	0.2
Maries Co	135	94.1	0.0	0.7	0.7	0.0	4.4
Marion Co	966	84.0	12.3	2.0	0.0	0.3	1.4
Mcdonald C	1,162	61.0	0.1	35.0	0.6	0.3	3.0
Mercer Co	127	93.7	0.0	1.6	0.0	0.0	4.7
Midtown WI	290	19.7	71.4	2.1	0.0	2.1	4.8
Miller co	673	92.1	1.5	3.6	0.6	0.4	1.8
Mississipp	788	59.4	38.2	1.5	0.3	0.3	0.4
Moniteau C	276	83.0	0.7	14.9	0.0	0.0	1.4
Monroe Co	296	81.8	14.5	2.7	0.0	0.3	0.7

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with errors.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 4A
2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Racial and Ethnic Distribution by Clinic						
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other / Unknown %
Montgomery	290	89.0	3.8	3.8	0.7	0.0	2.8
Morgan Co	575	94.4	1.2	1.4	0.2	0.2	2.6
NE Communi	629	82.5	7.5	7.5	0.2	0.3	2.1
New Madrid	745	63.1	34.6	2.0	0.0	0.0	0.3
Newton Co	1,602	84.1	0.9	10.0	1.0	2.1	1.8
Nmhc-WIC-A	673	94.4	1.2	1.9	0.0	1.0	1.5
Nodaway Co	366	96.4	0.8	0.5	0.0	0.0	2.2
North Coun	279	6.1	91.8	1.1	0.4	0.0	0.7
Oregon Co	598	97.0	0.2	2.2	0.0	0.0	0.7
Ozark Co H	283	95.1	0.0	2.5	0.4	0.4	1.8
Pemiscot C	1,256	49.6	47.1	2.8	0.0	0.1	0.5
Perry Co H	499	93.0	1.2	2.6	0.6	0.0	2.6
Pettis Cou	1,563	72.3	3.6	17.4	0.3	0.4	6.0
Phelps Co	1,205	86.8	1.8	2.0	0.5	1.6	7.3
Pike Co He	499	80.8	9.8	8.6	0.0	0.0	0.8
Pine Lawn	1,381	3.8	91.4	2.6	0.0	0.4	1.8
Platte Chd	1,280	66.6	11.4	12.3	0.4	4.0	5.4
Platte Co	173	80.3	1.7	12.7	0.0	0.0	5.2
Polk Co He	847	95.3	0.9	2.0	0.2	0.1	1.4

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with errors.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 4A
2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Racial and Ethnic Distribution by Clinic						
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other / Unknown %
Polo-Braym	77	*	*	*	*	*	*
Prince Hal	589	0.5	92.5	2.4	0.0	0.0	4.6
Pulaski Co	531	91.7	0.6	5.5	0.2	0.4	1.7
Putnam Co	165	98.2	0.0	1.2	0.0	0.6	0.0
Ralls Co H	144	93.1	3.5	2.1	0.0	0.7	0.7
Randolph C	1,032	86.1	6.7	2.5	0.4	0.9	3.4
Ray Co Hea	528	93.6	4.0	0.9	0.4	0.0	1.1
Reynolds C	238	96.6	0.8	1.7	0.0	0.4	0.4
Richard Ca	180	2.8	1.1	83.3	0.0	0.6	12.2
Ripley Co	560	93.9	2.0	2.5	0.2	0.2	1.3
Saline Co	666	73.0	7.2	12.5	0.0	0.9	6.5
Sam-Rod-Hi	180	89.4	3.9	3.9	0.0	0.0	2.8
Sam-Rod-Le	243	72.0	5.3	19.8	0.0	0.0	2.9
Sam-Rod-Od	183	90.7	3.3	1.6	0.5	0.0	3.8
Samuel U R	2,485	10.7	22.3	47.2	0.1	9.1	10.6
Schuyler C	124	88.7	0.0	8.9	0.8	0.0	1.6
Scotland C	96	*	*	*	*	*	*
Scott Co H	1,639	73.4	22.9	2.6	0.2	0.1	0.8
Sears Cert	1,427	2.0	92.4	2.9	0.1	0.6	2.0

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with errors.
*   Percentages are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 4A

2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years (2)

Page: 8
Run Date: 7 / 8 / 04

Comparison of Racial and Ethnic Distribution by Clinic						
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other / Unknown %
Shannon Co	390	96.9	0.0	2.3	0.5	0.0	0.3
Shelby Co	257	95.3	1.9	0.8	0.4	1.2	0.4
SO Central	1,504	96.0	0.5	1.9	0.5	0.1	1.1
Soulard/Gr	423	8.3	44.4	36.6	0.0	2.1	8.5
South Coun	1,903	72.2	9.2	7.4	0.2	4.4	6.7
South Grac	338	6.5	36.7	50.3	0.0	0.9	5.6
South Poin	1,591	22.3	56.5	4.7	0.3	6.0	10.2
Southside	87	*	*	*	*	*	*
Spgfld-Gre	6,104	84.7	4.6	6.1	0.3	1.5	2.8
St Charles	2,248	70.2	11.4	11.9	0.3	2.4	3.8
St Clair C	352	89.2	0.6	1.7	0.0	0.0	8.5
St Francoi	1,695	95.5	1.2	2.2	0.0	0.2	0.9
St Joseph-	1,927	80.4	8.3	3.5	0.4	0.1	7.3
St Louis C	2,273	43.6	46.8	3.9	0.1	1.5	4.1
St Louis H	3	*	*	*	*	*	*
St Lukes N	1,043	20.8	44.1	30.7	0.5	1.3	2.6
St Stephen	9	*	*	*	*	*	*
Starkloff-	569	6.9	83.1	3.3	0.2	2.3	4.2
StCHarvstr	113	69.9	13.3	7.1	1.8	1.8	6.2

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with errors.
 * Percentages are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 4A
2003 Pediatric Nutrition Surveillance [1]
Missouri
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years [2]

Comparison of Racial and Ethnic Distribution by Clinic						
Clinic	Number Included in Analysis [3]	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other / Unknown %
StChrlsVFW	306	84.3	6.9	2.9	0.7	1.6	3.6
Ste Genevi	369	96.2	1.9	1.4	0.0	0.3	0.3
Stoddard C	915	97.9	0.7	1.0	0.0	0.0	0.4
Stone Co H	626	97.9	0.2	1.3	0.0	0.2	0.5
Sullivan C	282	53.5	2.1	43.3	0.4	0.4	0.4
Swope Pkwy	2,421	7.0	86.5	3.9	0.3	0.5	1.8
Taney Coun	670	93.3	0.7	3.9	0.3	0.0	1.8
Texas Co H	785	95.7	1.1	1.7	0.1	0.3	1.1
Tmce-Grand	2,056	52.1	25.7	15.8	0.4	0.6	5.4
Tmce-Jacks	2,708	78.3	6.4	8.6	0.8	1.8	4.2
Tmce-Lee's	19	*	*	*	*	*	*
Tmce-Oak G	276	90.6	0.7	5.4	0.4	0.0	2.9
Tmce-Susqu	220	83.2	4.1	6.8	0.5	1.8	3.6
Tmce-Van H	79	*	*	*	*	*	*
Tmcw-Ldrp	979	11.7	61.0	22.7	0.1	1.4	3.1
Tmcw-OB Cl	594	14.6	56.4	22.7	0.0	1.3	4.9
Tmcw-Pedia	632	12.8	54.1	27.4	0.0	1.6	4.1
Tri-County	192	94.3	1.0	1.6	1.0	0.0	2.1
Vernon Co	661	76.7	0.8	2.6	0.0	0.0	20.0

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with errors.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 4A

2003 Pediatric Nutrition Surveillance (1)

Missouri

Comparison of Racial and Ethnic Distribution by Clinic

Children Aged < 5 Years (2)

Page: 10

Run Date: 7 / 8 / 04

Comparison of Racial and Ethnic Distribution by Clinic						
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other / Unknown %
Washington	923	95.6	1.5	1.8	0.3	0.0	0.8
Watertower	709	1.0	95.6	2.1	0.0	0.1	1.1
Wayne Co H	611	97.4	0.5	1.1	0.2	0.2	0.7
Webster Co	1,175	95.2	0.6	2.5	0.4	0.0	1.3
Wellston W	29	*	*	*	*	*	*
Wentzville	1	*	*	*	*	*	*
Worth-Tri	49	*	*	*	*	*	*
Wright Cou	759	96.6	0.5	1.8	0.5	0.3	0.3
Unknown Clinics	1,059	4.2	33.8	57.6	0.0	0.4	4.1
Missouri	**136,951**	**65.7**	**21.2**	**8.3**	**0.3**	**1.1**	**3.4**
Nation (Prior Year)	**5,333,226**	**39.1**	**20.8**	**32.2**	**1.2**	**3.3**	**3.5**

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with errors.
*   Percentages are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 5A
2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
(Delmar) P	1,556	41.5	23.7	34.8
(Maplewood	569	36.0	25.7	38.3
00Family C	1	*	*	*
00Miller C	127	35.4	25.2	39.4
01SL Conne	455	32.7	25.3	42.0
02Family C	552	30.6	28.8	40.6
Andrew Co	354	31.6	21.5	46.9
Atchison C	178	22.5	32.0	45.5
Audrain Co	576	37.7	24.7	37.7
Barnes Hos	168	100.0	0.0	0.0
Barry Co H	1,500	29.3	26.3	44.4
Barton Co	480	23.8	27.5	48.8
Bates Co H	525	30.1	25.0	45.0
BeltonCass	115	8.7	28.7	62.6
Benton Co	429	27.0	26.1	46.9
Blue Sprin	1,335	32.0	26.4	41.6
Bollinger	391	25.8	27.9	46.3
BransTaney	1,080	26.6	27.8	45.6

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
 * Percentages are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 5A
2003 Pediatric Nutrition Surveillance [1]
Missouri
Comparison of Age Distribution by Clinic
Children Aged < 5 Years [2]

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Butler Co	1,645	29.8	26.6	43.6
Callaway C	651	33.8	26.7	39.5
Camden Co	896	32.4	25.4	42.2
Cape Girar	1,361	35.6	26.7	37.8
Carondelet	1,244	34.4	26.5	39.1
Carroll Co	308	26.0	26.6	47.4
Carter Co	265	19.6	27.2	53.2
Cass Co He	1,164	32.8	22.6	44.6
Cedar Co H	482	26.3	22.2	51.5
Chariton C	151	27.8	29.1	43.0
Christian	1,442	26.8	26.8	46.5
Clark Co H	207	27.5	25.6	46.9
Clay Chd-E	61	*	*	*
Clay Chd-P	50	*	*	*
Clay Co He	2,344	37.1	23.8	39.0
Clay Metro	17	*	*	*
Clinton Co	573	29.8	24.1	46.1
Cole Co He	1,781	30.4	25.4	44.2

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
 * Percentages are not calculated if < 100 records are available for analysis after exclusions.

CDC

2003 Pediatric Nutrition Surveillance [1]
Missouri
Comparison of Age Distribution by Clinic
Children Aged < 5 Years [2]

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Columbia-B	2,055	35.4	25.8	38.7
Comp Healt	735	36.5	26.5	37.0
Cooper Co	339	33.9	26.5	39.5
Crawford C	718	31.1	24.5	44.4
Crescent C	1,503	31.4	28.3	40.3
Dade Co He	276	27.5	23.6	48.9
Dallas Co	410	29.8	28.3	42.0
Daviess Co	284	29.2	25.7	45.1
Dekalb Co-	166	27.1	22.9	50.0
Dent Co He	521	26.7	25.1	48.2
Douglas Co	418	28.5	24.6	46.9
Dunklin Co	1,082	29.9	26.4	43.7
Elsber-Lnc	51	*	*	*
Florence H	891	31.0	26.4	42.6
Forest Par	751	52.2	24.2	23.6
Franklin C	1,597	35.3	26.0	38.7
FtWood-Pul	1,064	29.3	25.6	45.1
Gasconade	366	28.7	27.6	43.7

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
 * Percentages are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 5A
2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
GascondOsg	161	26.1	24.2	49.7
Grace Hill	340	40.0	25.3	34.7
Grundy Co	362	27.9	24.6	47.5
Hamilton/C	111	24.3	22.5	53.2
Harrison C	297	29.0	24.6	46.5
Hartville-	152	23.0	27.0	50.0
Henry Co H	647	26.6	28.9	44.5
HGP-Connec	178	28.1	31.5	40.4
Hickory Co	193	33.7	23.8	42.5
Holt Count	103	35.9	24.3	39.8
Hosphill-T	2,767	15.6	35.2	49.2
Howard Co	175	32.0	25.1	42.9
Iron Co He	405	25.7	26.7	47.7
Jfcac-Arno	1,361	38.0	24.0	38.1
Jfcac-Unit	1,706	36.2	25.7	38.1
John C Mur	2,540	44.8	22.3	32.9
Johnson Co	1,423	31.5	26.9	41.6
Joplin Cit	4,015	31.0	26.3	42.7

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
 * Percentages are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 5A
2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Kansas Cit	2	*	*	*
Kchd-Ruski	2	*	*	*
Knox Co He	151	19.9	25.8	54.3
La Clinica	849	28.6	25.2	46.2
Laclede Co	1,017	34.6	25.6	39.8
Lakewood-T	1,638	38.0	26.8	35.2
Lawrence C	1,232	27.9	26.0	46.1
Lewis Co H	315	29.8	24.1	46.0
Lincoln Co	782	33.4	27.7	38.9
Linn Co He	360	28.3	25.0	46.7
Livingston	428	29.9	28.0	42.1
Macon Co H	408	27.7	24.0	48.3
Madison Co	546	26.6	25.8	47.6
Malden-Dnk	548	25.9	26.5	47.6
Maries Co	135	33.3	21.5	45.2
Marion Co	966	27.8	29.5	42.7
Mcdonald C	1,162	29.7	23.7	46.6
Mercer Co	127	29.1	29.9	40.9

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
 * Percentages are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 5A
2003 Pediatric Nutrition Surveillance [1]
Missouri
Comparison of Age Distribution by Clinic
Children Aged < 5 Years [2]

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Midtown WI	290	34.1	25.2	40.7
Miller co	673	29.9	23.0	47.1
Mississipp	788	21.7	26.8	51.5
Moniteau C	276	30.8	21.4	47.8
Monroe Co	296	24.3	22.3	53.4
Montgomery	290	29.7	24.8	45.5
Morgan Co	575	27.5	26.4	46.1
NE Communi	629	30.2	26.7	43.1
New Madrid	745	27.8	22.6	49.7
Newton Co	1,602	26.7	27.8	45.6
Nmhc-WIC-A	673	27.9	27.5	44.6
Nodaway Co	366	31.1	23.0	45.9
North Coun	279	44.8	20.4	34.8
Oregon Co	598	21.9	23.1	55.0
Ozark Co H	283	25.1	28.6	46.3
Pemiscot C	1,256	25.6	27.5	46.8
Perry Co H	499	27.5	24.6	47.9
Pettis Cou	1,563	28.1	25.3	46.6

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
 * Percentages are not calculated if < 100 records are available for analysis after exclusions.

CDC

2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Phelps Co	1,205	30.3	27.6	42.2
Pike Co He	499	31.9	29.3	38.9
Pine Lawn	1,381	41.1	22.8	36.1
Platte Chd	1,280	29.2	28.8	42.0
Platte Co	173	26.6	34.1	39.3
Polk Co He	847	27.0	24.4	48.5
Polo-Braym	77	*	*	*
Prince Hal	589	28.4	25.0	46.7
Pulaski Co	531	30.9	25.2	43.9
Putnam Co	165	21.2	28.5	50.3
Ralls Co H	144	34.0	18.8	47.2
Randolph C	1,032	25.3	26.2	48.5
Ray Co Hea	528	30.9	25.0	44.1
Reynolds C	238	25.6	29.0	45.4
Richard Ca	180	32.8	26.7	40.6
Ripley Co	560	23.0	26.6	50.4
Saline Co	666	28.8	28.1	43.1
Sam-Rod-Hi	180	28.9	22.2	48.9

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
 * Percentages are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 5A
2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Sam-Rod-Le	243	28.0	18.9	53.1
Sam-Rod-Od	183	27.3	28.4	44.3
Samuel U R	2,485	26.3	25.7	48.0
Schuyler C	124	21.8	31.5	46.8
Scotland C	96	*	*	*
Scott Co H	1,639	29.6	25.3	45.1
Sears Cert	1,427	29.4	25.1	45.6
Shannon Co	390	22.1	27.9	50.0
Shelby Co	257	24.5	25.7	49.8
SO Central	1,504	29.3	26.3	44.5
Soulard/Gr	423	33.1	27.2	39.7
South Coun	1,903	44.5	24.3	31.2
South Grac	338	54.4	25.7	19.8
South Poin	1,591	28.7	27.6	43.7
Southside	87	*	*	*
Spgfld-Gre	6,104	32.9	25.6	41.5
St Charles	2,248	39.9	24.7	35.4
St Clair C	352	21.0	27.3	51.7

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
 * Percentages are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 5A
2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
St Francoi	1,695	32.2	26.5	41.3
St Joseph-	1,927	36.4	25.4	38.1
St Louis C	2,273	46.9	23.0	30.1
St Louis H	3	*	*	*
St Lukes N	1,043	38.6	25.1	36.2
St Stephen	9	*	*	*
Starkloff-	569	38.8	26.2	35.0
StCHarvstr	113	33.6	26.5	39.8
StChrlsVFW	306	36.6	21.9	41.5
Ste Genevi	369	29.3	24.9	45.8
Stoddard C	915	27.7	29.1	43.3
Stone Co H	626	30.4	25.2	44.4
Sullivan C	282	30.5	24.5	45.0
Swope Pkwy	2,421	35.6	26.4	38.1
Taney Coun	670	23.9	27.2	49.0
Texas Co H	785	26.0	26.4	47.6
Tmce-Grand	2,056	30.3	27.0	42.8
Tmce-Jacks	2,708	29.0	27.9	43.2

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
 * Percentages are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 5A
2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Tmce-Lee's	19	*	*	*
Tmce-Oak G	276	30.8	26.1	43.1
Tmce-Susqu	220	22.7	26.4	50.9
Tmce-Van H	79	*	*	*
Tmcw-Ldrp	979	90.0	5.0	5.0
Tmcw-OB Cl	594	37.9	22.7	39.4
Tmcw-Pedia	632	17.6	28.0	54.4
Tri-County	192	30.7	23.4	45.8
Vernon Co	661	28.4	28.0	43.6
Washington	923	30.6	27.4	42.0
Watertower	709	42.6	23.8	33.6
Wayne Co H	611	23.1	24.7	52.2
Webster Co	1,175	27.5	24.2	48.3
Wellston W	29	*	*	*
Wentzville	1	*	*	*
Worth-Tri	49	*	*	*
Wright Cou	759	24.2	27.7	48.1
Unknown Clinics	1,059	37.1	26.4	36.4

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
 * Percentages are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 5A
2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Missouri	**136,951**	**32.1**	**25.8**	**42.1**
Nation (Prior Year)	**5,333,226**	**36.6**	**21.7**	**41.7**

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
 * Percentages are not calculated if < 100 records are available for analysis after exclusions.

CDC

2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)

Clinic	Birthweight (5) Low Number	% <2500 g (Rank)	High % >4000 g (Rank)	Height and Weight Number	Short Stature (6) % <5th (Rank)	Under-weight (7) % <5th (Rank)	Over-weight (7) % >95th (Rank)	≥ 2 Yrs Over-weight (7) Number	% 85th- <95th (Rank)	% >95th (Rank)	Anemia Low Hb/Hct (8) Number	% (Rank)
(Delmar) P	611	13.1 (101)	3.6 (7)	1,496	8.2 (13	9.1 (14	9.7 (26)	532	17.5 (83)	16.7 (10	847	29.5 (146)
(Maplewood	184	12.5 (99)	7.6 (72)	529	9.8 (15	7.4 (13	18.0 (15	211	19.0 (11	25.6 (13	338	19.5 (119)
00Family C	0	* *	* *	1	* *	* *	* *	1	* *	* *	1	* *
00Miller C	38	* *	* *	123	6.5 (73)	2.4 (39)	8.1 (12	52	* *	* *	83	* *
01SL Conne	140	12.1 (96)	6.4 (44)	444	8.8 (14	8.1 (14	17.8 (15	185	22.7 (12	19.5 (12	286	10.5 (43)
02Family C	160	16.9 (112)	5.6 (36)	532	7.9 (12	10.0 (15	11.8 (69)	220	15.5 (54)	11.4 (36)	357	23.0 (132)
Andrew Co	102	4.9 (6)	6.9 (56)	347	4.9 (23)	5.8 (12	11.5 (60)	171	12.3 (9)	12.9 (62)	222	4.5 (5)
Atchison C	32	* *	* *	175	8.6 (13	1.7 (15	19.4 (15	85	* *	* *	134	23.9 (136)
Audrain Co	191	7.9 (34)	5.2 (30)	561	13.2 (16	2.1 (26)	15.9 (13	217	15.2 (46)	16.1 (10	357	21.3 (127)
Barnes Hos	166	12.0 (91)	7.8 (77)	161	5.0 (26)	14.9 (16	1.9 (1)	0	* *	* *	1	* *
Barry Co H	393	6.6 (16)	8.4 (92)	1,461	5.1 (30)	3.1 (72)	11.9 (72)	638	12.2 (8)	13.5 (68)	989	11.6 (56)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 6A
2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 2
Run Date: 7 / 8 / 04

Comparison of Growth and Anemia Indicators by Clinic (3,4)

Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature (6)	Under-weight (7)	Over-weight (7)		≥ 2 Yrs Over-weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- <95th (Rank)	% >95th (Rank)	Number	% (Rank)
Barton Co	105	4.8 (3)	10.5 (109)	470	6.0 (57)	3.0 (68)	12.8 (94)	230	23.9 (13	15.2 (97)	338	15.4 (84)
Bates Co H	135	7.4 (26)	5.9 (39)	504	7.9 (12	5.2 (11	11.9 (72)	226	23.0 (13	18.1 (11	336	19.6 (120)
BeltonCass	10	* *	* *	115	2.6 (2)	3.5 (83)	13.0 (10	68	* *	* *	92	* *
Benton Co	107	8.4 (42)	6.5 (45)	417	9.1 (14	6.2 (13	10.6 (45)	201	17.9 (91)	15.9 (10	293	12.6 (66)
Blue Sprin	366	10.9 (87)	6.8 (51)	1,291	6.0 (57)	4.0 (98)	9.1 (19)	550	18.0 (95)	9.1 (11)	834	12.2 (62)
Bollinger	93	* *	* *	379	5.0 (26)	2.4 (39)	10.3 (39)	177	20.3 (12	11.9 (47)	267	7.9 (19)
BransTaney	259	10.0 (68)	7.3 (65)	1,039	6.4 (67)	1.6 (13)	11.4 (58)	489	18.6 (10	10.8 (25)	725	10.5 (43)
Butler Co	436	9.9 (65)	3.9 (12)	1,587	11.4 (15	1.9 (20)	16.4 (14	686	20.8 (12	18.1 (11	1,074	12.1 (60)
Callaway C	200	10.0 (68)	7.5 (70)	625	9.4 (14	2.1 (26)	11.2 (55)	250	15.2 (46)	10.8 (25)	416	11.3 (51)
Camden Co	249	4.8 (3)	7.6 (72)	874	4.6 (18)	1.7 (15)	8.9 (17)	373	14.2 (32)	8.3 (7)	565	9.4 (33)
Cape Girar	421	9.3 (57)	6.9 (56)	1,301	7.6 (10	3.3 (76)	12.2 (78)	498	16.5 (69)	12.0 (50)	842	16.9 (97)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 6A
2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 3
Run Date: 7 / 8 / 04

	Comparison of Growth and Anemia Indicators by Clinic (3,4)										
	Birthweight (5)		Height and Weight							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under-weight (7)	Over-weight (7)	≥ 2 Yrs Over-weight (7)				
Clinic	Number / % <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- <95th (Rank)	% >95th (Rank)	Number	% (Rank)
Carondelet	384 10.4 (78)	8.1 (85)	1,208	7.6(10	8.2(14	11.5(60)	477	14.0(28)	14.3(83)	806	10.0 (40)
Carroll Co	74 * *	* *	296	4.7(21	2.4(39	11.1(54)	145	16.6(71)	10.3(23)	204	12.7 (70)
Carter Co	48 * *	* *	258	3.5 (6)	7.4(13	5.4 (3)	141	8.5 (1)	7.1 (4)	186	17.2(101)
Cass Co He	317 7.9 (34)	6.9 (56)	1,122	6.9(87)	3.6(87)	10.1(36)	493	14.6(36)	9.1(11)	736	18.9(114)
Cedar Co H	115 8.7 (46)	10.4 (108)	467	4.5(16)	3.6(87)	15.8(13	242	19.8(11	18.6(11	330	20.0(122)
Chariton C	36 * *	* *	143	3.5 (6)	0.0 (1)	12.6(87)	62	* *	* *	106	3.8 (2)
Christian	338 6.8 (21)	6.8 (51)	1,408	5.5(41)	4.4(10	9.4(21)	662	14.0(28)	10.1(20)	992	11.6 (56)
Clark Co H	52 * *	* *	202	5.0(26)	3.0(68)	9.9(32)	96	* *	* *	143	7.0 (13)
Clay Chd-E	5 * *	* *	58	* *	* *	* *	38	* *	* *	44	* *
Clay Chd-P	12 * *	* *	49	* *	* *	* *	23	* *	* *	35	* *
Clay Co He	759 7.5 (30)	8.8 (96)	2,265	7.7(11	4.7(11	12.5(85)	879	17.9(91)	13.5(68)	1,397	18.9(114)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

CDC

2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)										
	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)			
Clinic	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- <95th (Rank)	% >95th (Rank)	Number	% (Rank)
Clay Metro	2	* *	* *	16	* *	* *	* *	11	* *	* *	12	* *
Clinton Co	155	10.3 (76)	11.0 (110)	561	6.6 (76)	3.0 (68)	13.7 (10)	261	20.7 (12	17.2 (11	402	8.0 (20)
Cole Co He	475	8.8 (48)	9.9 (106)	1,694	5.5 (41)	4.3 (10	10.4 (40)	764	16.4 (65)	11.6 (44)	1,182	3.8 (2)
Columbia-B	631	9.8 (63)	7.0 (61)	1,958	6.9 (87)	3.2 (75)	11.8 (69)	774	16.9 (75)	14.1 (81)	1,272	24.8 (139)
Comp Healt	252	10.3 (76)	2.8 (4)	709	5.4 (39)	8.6 (14	9.7 (26)	270	13.7 (24)	14.8 (92)	458	18.6 (113)
Cooper Co	105	14.3 (108)	4.8 (26)	331	5.1 (30)	2.4 (39)	12.1 (75)	134	26.1 (13	14.9 (94)	207	12.6 (66)
Crawford C	207	7.2 (24)	5.3 (33)	704	5.8 (52)	3.3 (76)	10.2 (38)	316	14.9 (41)	14.6 (89)	461	16.9 (97)
Crescent C	415	9.4 (58)	8.0 (83)	1,455	5.8 (52)	5.9 (12	10.9 (50)	588	14.1 (30)	9.4 (14	991	33.8 (150)
Dade Co He	70	* *	* *	267	7.9 (12	5.2 (11	9.7 (26)	134	13.4 (16	13.4 (67)	172	8.7 (27)
Dallas Co	120	10.0 (68)	9.2 (101)	391	8.4 (13	3.3 (76)	17.1 (14	162	13.6 (22)	14.2 (82)	233	18.0 (109)
Daviess Co	71	* *	* *	276	10.9 (15	1.1 (6)	17.8 (15	125	18.4 (10	19.2 (12	194	4.1 (4)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

	Comparison of Growth and Anemia Indicators by Clinic (3,4)										
	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)
	Low	High		Short Stature (6)	Under-weight (7)	Over-weight (7)		≥ 2 Yrs Over-weight (7)			
Clinic	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Dekalb Co-	36	* *	* *	166	7.2(10	2.4(39)	13.3(10	83	* *	* *	120	9.2 (32)
Dent Co He	117	4.3 (1)	6.8 (51)	507	8.7(13	2.0(22)	9.3(20)	249	14.9(41)	6.8 (3)	354	10.7 (46)
Douglas Co	103	8.7 (46)	4.9 (28)	409	4.6(18)	5.9(12	10.5(44)	199	18.1(99)	9.0(10)	274	13.9 (75)
Dunklin Co	285	8.8 (48)	3.9 (12)	1,001	7.3(10	2.0(22)	16.1(14	437	18.5(10	19.7(12	693	11.1 (50)
Elsber-Lnc	10	* *	* *	49	* *	* *	* *	23	* *	* *	36	* *
Florence H	262	16.0(111)	4.2 (16)	851	7.6(10	9.4(14	13.5(10	370	15.1(45)	18.1(11	531	10.9 (49)
Forest Par	383	9.1 (51)	7.3 (65)	722	5.1(30)	9.8(14	10.9(50)	174	17.2(79)	19.0(12	336	28.3 (144)
Franklin C	504	8.1 (38)	8.9 (97)	1,567	7.5(10	4.0(98)	11.5(60)	613	14.5(35)	11.9(47)	990	9.6 (38)
FtWood-Pul	270	6.7 (19)	10.0(107)	954	6.1(61)	2.5(47)	13.7(10	458	10.9 (4)	8.1 (6)	651	34.1 (151)
Gasconade	100	12.0 (91)	4.0 (15)	356	5.1(30)	3.1(72)	11.0(53)	160	19.4(11	15.0(95)	245	6.5 (11)
GascondOsg	40	* *	* *	153	5.9(55)	4.6(10	15.7(13	74	* *	* *	109	15.6 (86)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 6A
2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low	High			Short Stature (6)	Under-weight (7)	Over-weight (7)		≥ 2 Yrs Over-weight (7)			
Clinic	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- <95th (Rank)	% >95th (Rank)	Number	% (Rank)
Grace Hill	129	10.1 (74)	4.7 (25)	319	7.8(11	12.2(15	11.6(67)	109	12.8(12)	17.4(11	180	16.7 (94)
Grundy Co	88	* *	* *	348	5.5(41)	4.9(11	11.5(60)	163	17.2(79)	12.9(62)	251	8.4 (25)
Hamilton/C	25	* *	* *	108	6.5(73)	0.9 (5)	16.7(14	60	* *	* *	80	* *
Harrison C	78	* *	* *	286	4.2(13)	3.1(72)	16.1(14	133	13.5(19)	20.3(12	202	12.9 (71)
Hartville-	34	* *	* *	149	3.4 (5)	6.0(12	6.7 (6)	75	* *	* *	105	23.8 (135)
Henry Co H	155	8.4 (42)	12.9 (111)	637	7.1(95)	2.4(39)	14.0(11	285	18.2(10	13.7(73)	443	17.8 (107)
HGP-Connec	48	* *	* *	173	10.4(15	8.1(14	10.4(40)	72	* *	* *	118	16.1 (89)
Hickory Co	63	* *	* *	188	4.8(22)	2.7(50)	14.4(12	82	* *	* *	122	8.2 (22)
Holt Count	30	* *	* *	100	5.0(26)	0.0 (1)	14.0(11	41	* *	* *	66	* *
Hosphill-T	370	12.2 (97)	5.7 (38)	2,669	8.3(13	4.0(98)	15.2(13	1,353	17.8(88)	14.0(79)	2,194	25.0 (140)
Howard Co	50	* *	* *	168	3.6 (8)	2.4(39)	10.1(36)	73	* *	* *	106	9.4 (33)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)

Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Number	Low % <2500 g (Rank)	High % >4000 g (Rank)	Number	Short Stature (6) % <5th (Rank)	Under-weight (7) % <5th (Rank)	Over-weight (7) % >95th (Rank)	≥ 2 Yrs Over-weight (7) Number	% 85th- <95th (Rank)	% >95th (Rank)	Number	% (Rank)
Iron Co He	99	* *	* *	394	7.1 (95)	2.0 (22)	14.7 (12	189	19.0 (11	14.8 (92)	283	7.4 (16)
Jfcac-Arno	459	7.4 (26)	8.1 (85)	1,342	5.6 (45)	3.5 (83)	9.5 (23)	501	15.4 (53)	13.8 (75)	803	14.2 (76)
Jfcac-Unit	544	8.6 (45)	7.2 (64)	1,658	6.9 (87)	4.5 (10	8.6 (15)	634	13.7 (24)	12.8 (61)	1,046	17.0 (100)
John C Mur	1,050	10.4 (78)	4.4 (21)	2,433	7.1 (95)	10.0 (15	8.1 (12)	808	12.1 (7)	10.0 (18)	1,379	20.2 (124)
Johnson Co	398	4.8 (3)	6.0 (41)	1,374	6.9 (87)	2.3 (33)	12.4 (83)	570	14.2 (32)	11.9 (47)	906	17.7 (105)
Joplin Cit	1,071	9.2 (55)	7.8 (77)	3,865	7.3 (10	2.7 (50)	12.2 (78)	1,677	17.7 (86)	13.5 (68)	2,408	18.1 (110)
Kansas Cit	0	* *	* *	2	* *	* *	* *	0	* *	* *	2	* *
Kchd-Ruski	0	* *	* *	2	* *	* *	* *	1	* *	* *	1	* *
Knox Co He	24	* *	* *	149	4.0 (11)	2.0 (22)	13.4 (10	80	* *	* *	107	5.6 (6)
La Clinica	222	9.9 (65)	6.8 (51)	824	6.7 (81)	10.2 (15	14.2 (11	378	13.8 (26)	18.0 (11	582	23.2 (133)
Laclede Co	316	8.5 (44)	8.2 (87)	988	7.5 (10	5.1 (11	9.8 (29)	402	16.4 (65)	10.9 (30)	632	13.6 (74)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 6A
2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature (6)	Under-weight (7)	Over-weight (7)		≥ 2 Yrs Over-weight (7)			
Clinic	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Lakewood-T	548	8.8 (48)	8.6 (93)	1,590	7.0(92)	5.6(12	12.9(99)	563	17.9(91)	12.1(52)	974	19.6(120)
Lawrence C	301	6.6 (16)	9.0 (99)	1,197	5.8(52)	6.1(12	10.4(40)	554	12.5(11)	10.8(25)	815	15.1 (81)
Lewis Co H	73	* *	* *	304	3.9 (9)	3.6(87)	12.8(94)	146	16.4(65)	16.4(10	205	10.7 (46)
Lincoln Co	231	7.4 (26)	7.4 (68)	747	7.4(10	2.8(54)	14.3(11	292	17.8(88)	15.4(99)	496	12.9 (71)
Linn Co He	100	12.0 (91)	6.0 (41)	351	5.7(48)	0.6 (3)	18.5(15	168	18.5(10	23.2(13	243	11.9 (59)
Livingston	110	6.4 (13)	3.6 (7)	420	7.1(95)	3.3(76)	12.6(87)	182	14.8(40)	13.7(73)	282	7.4 (16)
Macon Co H	104	10.6 (84)	8.7 (94)	401	8.2(13	2.2(32)	17.7(15	190	20.0(11	21.1(12	270	9.6 (38)
Madison Co	135	7.4 (26)	5.2 (30)	529	6.4(67)	4.5(10	18.0(15	256	18.0(95)	25.0(13	365	8.2 (22)
Malden-Dnk	129	13.2 (102)	3.9 (12)	528	7.6(10	2.3(33)	16.3(14	252	20.6(12	17.1(10	375	12.5 (65)
Maries Co	39	* *	* *	131	3.1 (4)	6.9(13	15.3(13	62	* *	* *	87	* *
Marion Co	251	13.5 (104)	5.2 (30)	910	9.6(14	2.9(57)	15.2(13	403	20.1(11	11.4(36)	610	15.9 (87)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 6A
2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)
	Low	High		Short Stature (6)	Under-weight (7)	Over-weight (7)		≥ 2 Yrs Over-weight (7)			
Clinic	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Mcdonald C	315	6.7 (19)	7.6 (72)	1,083	9.1(14	2.1 (26)	19.1(15	501	15.8 (60)	16.6(10	716	12.6 (66)
Mercer Co	36	* *	* *	122	4.9(23)	4.1(10	13.9(11	49	* *	* *	80	* *
Midtown WI	93	* *	* *	280	6.8(83)	10.4(15	7.5(10)	116	17.2(79)	11.2(33)	179	28.5 (145)
Miller co	186	8.1 (38)	5.9 (39)	653	8.7(13	1.7(15)	14.1(11	305	18.4(10	12.1(52)	446	18.4 (112)
Mississipp	158	10.1 (74)	3.8 (9)	757	8.7(13	2.1 (26)	12.9(99)	400	15.3(49)	14.3(83)	589	12.1 (60)
Moniteau C	71	* *	* *	275	7.6(10	2.9(57)	9.8(29)	127	20.5(12	11.8(45)	175	10.3 (41)
Monroe Co	66	* *	* *	289	7.6(10	2.1 (26)	14.5(12	155	16.8(73)	16.8(10	206	10.7 (46)
Montgomery	77	* *	* *	281	8.2(13	5.0(11	13.2(10	129	10.1 (2)	12.4(57)	192	7.3 (15)
Morgan Co	138	4.3 (1)	8.7 (94)	557	6.1(61)	2.9(57)	12.2(78)	264	15.2(46)	11.4(36)	386	21.0 (126)
NE Communi	174	9.8 (63)	6.9 (56)	600	5.5(41)	1.8(19)	11.5(60)	261	14.6(36)	17.6(11	406	11.3 (51)
New Madrid	181	13.3 (103)	1.7 (1)	724	9.3(14	2.9(57)	15.9(13	362	18.0(95)	15.2(97)	511	22.5 (129)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 6A
2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 10
Run Date: 7 / 8 / 04

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)
	Low	High			Short Stature (6)	Under-weight (7)	Over-weight (7)	≥ 2 Yrs Over-weight (7)			
Clinic	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- <95th (Rank)	% >95th (Rank)	Number	% (Rank)
Newton Co	372	5.1 (7)	9.1 (100)	1,551	3.9 (9)	4.6(10	9.0(18)	720	12.4(10)	10.4(24)	1,071	11.4 (53)
Nmhc-WIC-A	166	9.6 (61)	5.4 (34)	657	6.2(64)	4.9(11	8.2(14)	295	13.9(27)	11.5(41)	457	20.1 (123)
Nodaway Co	103	7.8 (33)	7.8 (77)	355	5.9(55)	3.9(94)	10.7(47)	167	13.2(15)	9.6(15)	236	15.3 (83)
North Coun	115	14.8 (109)	4.3 (17)	266	7.1(95)	8.3(14	11.7(68)	97	* *	* *	152	26.3 (141)
Oregon Co	114	10.5 (81)	14.0 (112)	566	2.5 (1)	3.7(90)	11.5(60)	316	18.4(10	12.0(50)	416	9.1 (30)
Ozark Co H	60	* *	* *	271	7.0(92)	1.5(11)	14.4(12	127	16.5(69)	15.0(95)	204	8.8 (28)
Pemiscot C	288	11.8 (90)	2.1 (3)	1,200	7.8(11	1.7(15)	15.2(13	578	19.7(11	17.0(10	876	24.0 (138)
Perry Co H	127	7.9 (34)	9.4 (105)	484	6.0(57)	2.7(50)	12.4(83)	235	21.7(12	14.0(79)	339	13.3 (73)
Pettis Cou	380	10.0 (68)	8.2 (87)	1,511	6.4(67)	2.3(33)	12.6(87)	708	17.2(79)	14.3(83)	1,031	10.3 (41)
Phelps Co	328	10.4 (78)	7.9 (81)	1,167	9.4(14	4.0(98)	12.7(91)	485	11.3 (6)	11.5(41)	816	11.6 (56)
Pike Co He	140	6.4 (13)	9.3 (102)	482	6.8(83)	3.9(94)	13.5(10	192	17.7(86)	13.0(64)	315	16.8 (95)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)

Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low		High		Short Stature (6)	Under-weight (7)	Over-weight (7)		≥ 2 Yrs Over-weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- <95th (Rank)	% >95th (Rank)	Number	% (Rank)
Pine Lawn	534	11.2 (89)	5.1 (29)	1,331	6.3 (66)	9.8 (14	9.5 (23)	488	13.5 (19)	12.1 (52)	798	14.9 (80)
Platte Chd	333	9.6 (61)	9.3 (102)	1,227	6.8 (83)	3.3 (76)	14.7 (12	523	15.3 (49)	14.5 (87)	802	19.0 (116)
Platte Co	43	* *	* *	165	7.3 (10	2.4 (39)	19.4 (15	70	* *	* *	120	14.2 (76)
Polk Co He	212	6.1 (11)	6.6 (47)	827	6.5 (73)	3.4 (81)	9.9 (32)	407	19.4 (11	9.6 (15)	574	10.6 (45)
Polo-Braym	18	* *	* *	74	* *	* *	* *	32	* *	* *	55	* *
Prince Hal	155	14.2 (107)	4.5 (22)	567	6.9 (87)	7.1 (13	12.0 (74)	269	16.0 (61)	10.8 (25)	398	23.9 (136)
Pulaski Co	141	9.9 (65)	3.5 (6)	505	9.9 (15	1.2 (9)	14.9 (12	228	16.7 (72)	13.2 (66)	333	23.4 (134)
Putnam Co	33	* *	* *	157	5.1 (30)	3.8 (93)	14.6 (12	82	* *	* *	118	3.4 (1)
Ralls Co H	42	* *	* *	137	8.0 (12	1.5 (11)	22.6 (16	68	* *	* *	90	* *
Randolph C	239	6.3 (12)	7.1 (62)	1,017	5.3 (36)	2.9 (57)	11.3 (57)	499	23.4 (13	14.6 (89)	711	18.3 (111)
Ray Co Hea	151	13.9 (106)	4.6 (23)	514	8.4 (13	3.9 (94)	7.2 (7)	225	18.2 (10	6.2 (1)	340	16.2 (92)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 6A
2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)										
	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)
	Low	High		Short Stature (6)	Under-weight (7)	Over-weight (7)	≥ 2 Yrs Over-weight (7)			
Clinic	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- <95th (Rank)	% >95th (Rank)	Number	% (Rank)
Reynolds C	55	* *	* *	233	5.6 (45)	2.6 (48)	15.9 (13	103	21.4 (12	19.4 (12	163	8.0 (20)
Richard Ca	31	* *	* *	174	5.7 (48)	1.1 (6)	15.5 (13	74	* *	* *	114	7.0 (13)
Ripley Co	115	12.2 (97)	1.7 (1)	549	8.6 (13	5.6 (12	8.6 (15)	275	14.9 (41)	8.0 (5)	402	12.4 (63)
Saline Co	174	5.2 (8)	4.6 (23)	650	6.6 (76)	2.3 (33)	12.8 (94)	282	15.6 (55)	12.1 (52)	442	17.4 (102)
Sam-Rod-Hi	46	* *	* *	175	5.7 (48)	1.1 (6)	17.1 (14	84	* *	* *	111	12.6 (66)
Sam-Rod-Le	60	* *	* *	231	7.8 (11	3.5 (83)	17.7 (15	127	19.7 (11	18.9 (12	155	16.1 (89)
Sam-Rod-Od	43	* *	* *	179	11.2 (15	0.6 (3)	12.8 (94)	82	* *	* *	131	17.6 (104)
Samuel U R	584	7.0 (22)	7.5 (70)	2,379	4.0 (11	5.6 (12	12.8 (94)	1,164	13.1 (13	13.1 (65)	1,707	19.4 (118)
Schuyler C	25	* *	* *	124	13.7 (16	7.3 (13	12.1 (75)	58	* *	* *	93	* *
Scotland C	27	* *	* *	94	* *	* *	* *	40	* *	* *	67	* *
Scott Co H	443	12.0 (91)	4.3 (17)	1,582	8.0 (12	2.1 (26)	10.6 (45)	718	16.0 (61)	12.5 (59)	1,075	16.6 (93)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 6A
2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)

Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature (6)	Under-weight (7)	Over-weight (7)		≥ 2 Yrs Over-weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- <95th (Rank)	% >95th (Rank)	Number	% (Rank)
Sears Cert	391	12.0 (91)	3.8 (9)	1,384	5.3 (36)	8.8 (14	9.8 (29)	634	15.6 (55)	11.5 (41)	951	30.1 (147)
Shannon Co	80	* *	* *	385	8.1 (12	2.6 (48)	14.0 (11	202	13.4 (16)	11.4 (36)	278	9.4 (33)
Shelby Co	54	* *	* *	245	4.5 (16)	2.9 (57)	12.7 (91)	127	15.7 (58)	10.2 (21)	180	9.4 (33)
SO Central	394	5.8 (9)	8.9 (97)	1,482	4.3 (15)	2.9 (57)	12.1 (75)	667	16.8 (73)	9.6 (15)	995	5.8 (7)
Soulard/Gr	130	10.0 (68)	7.7 (76)	394	11.2 (15	10.7 (15	20.8 (16	153	17.6 (85)	26.1 (13	259	6.6 (12)
South Coun	756	8.1 (38)	7.9 (81)	1,818	5.2 (35)	10.1 (15	9.4 (21)	586	15.0 (44)	15.7 (10	1,023	17.7 (105)
South Grac	177	10.7 (85)	6.2 (43)	320	9.1 (14	11.3 (15	15.0 (13	61	* *	* *	125	16.8 (95)
South Poin	422	9.2 (55)	6.9 (56)	1,533	5.3 (36)	7.3 (13	12.3 (82)	680	15.3 (49)	14.7 (91)	1,098	30.1 (147)
Southside	20	* *	* *	84	* *	* *	* *	42	* *	* *	62	* *
Spgfld-Gre	1,797	7.9 (34)	8.3 (89)	5,911	8.1 (12	3.0 (68)	13.3 (10	2,481	17.9 (91)	13.5 (68)	3,802	17.4 (102)
St Charles	776	7.5 (30)	7.6 (72)	1,972	6.6 (76)	3.7 (90)	11.5 (60)	764	15.3 (49)	13.5 (68)	1,168	8.2 (22)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 6A
2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature (6)	Under-weight (7)	Over-weight (7)	≥ 2 Yrs Over-weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- <95th (Rank)	% >95th (Rank)	Number	% (Rank)
St Clair C	65	* *	* *	333	7.8(11	3.9(94)	12.6(87)	172	19.8(11	12.2(56)	252	15.9 (87)
St Francoi	487	7.6 (32)	7.4 (68)	1,653	6.8(83)	2.9(57)	12.7(91)	684	18.1(99)	15.5(10	1,109	12.4 (63)
St Joseph-	635	9.1 (51)	6.5 (45)	1,852	6.7(81)	5.3(11	10.7(47)	705	16.9(75)	13.9(76)	1,163	14.4 (79)
St Louis C	992	10.7 (85)	6.7 (48)	2,190	4.9(23)	10.8(15	7.4 (8)	674	15.6(55)	11.4(36)	1,172	26.6 (142)
St Louis H	0	* *	* *	3	* *	* *	* *	2	* *	* *	3	* *
St Lukes N	374	9.1 (51)	8.3 (89)	1,001	7.7(11	6.1(12	13.4(10	374	14.7(38)	11.0(31)	567	15.5 (85)
St Stephen	4	* *	* *	9	* *	* *	* *	4	* *	* *	4	* *
Starkloff-	210	10.5 (81)	2.9 (5)	550	5.6(45)	12.7(16	6.5 (5)	194	13.4(16)	6.7 (2)	318	9.1 (30)
StCHarvstr	37	* *	* *	109	6.4(67)	2.8(54)	6.4 (4)	47	* *	* *	56	* *
StChrlsVFW	92	* *	* *	297	5.7(48)	4.4(10	10.4(40)	126	13.5(19)	8.7 (9)	164	8.5 (26)
Ste Genevi	96	* *	* *	364	9.6(14	1.4(10)	7.4 (8)	166	16.3(64)	10.2(21)	241	6.2 (9)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)
		Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)			
Clinic	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- <95th (Rank)	% >95th (Rank)	Number	% (Rank)
Stoddard C	232	9.1 (51)	5.6 (36)	881	7.5(10	2.3(33)	11.2(55)	390	21.0(12	14.4(86)	606	21.6 (128)
Stone Co H	170	7.1 (23)	7.1 (62)	614	4.2(13)	3.7(90)	7.7(11)	275	13.1(13)	8.4 (8)	399	11.5 (55)
Sullivan C	69	* *	* *	257	5.4(39)	2.3(33)	16.3(14	122	18.9(10	18.0(11	166	6.0 (8)
Swope Pkwy	797	9.5 (59)	5.5 (35)	2,336	7.6(10	6.5(13	9.5(23)	903	14.1(30)	10.0(18)	1,518	20.4 (125)
Taney Coun	138	6.5 (15)	8.0 (83)	647	6.6(76)	2.9(57)	10.0(34)	324	13.6(22)	9.3(13)	472	16.1 (89)
Texas Co H	180	8.3 (41)	6.7 (48)	756	4.6(18)	2.8(54)	10.8(49)	352	10.2 (3)	10.8(25)	531	9.0 (29)
Tmce-Grand	538	10.0 (68)	9.3 (102)	1,987	6.2(64)	4.6(10	11.8(69)	851	14.7(38)	11.8(45)	1,353	19.3 (117)
Tmce-Jacks	671	7.3 (25)	7.3 (65)	2,637	7.0(92)	3.5(83)	12.2(78)	1,145	17.5(83)	11.2(33)	1,800	14.3 (78)
Tmce-Lee's	0	* *	* *	19	* *	* *	* *	12	* *	* *	15	* *
Tmce-Oak G	72	* *	* *	270	7.8(11	5.9(12	16.3(14	117	11.1 (5)	14.5(87)	179	22.9 (131)
Tmce-Susqu	43	* *	* *	215	10.2(15	1.9(20)	14.0(11	111	18.0(95)	12.6(60)	155	30.3 (149)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 16
Run Date: 7 / 8 / 04

	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low		High		Short Stature (6)	Under-weight (7)	Over-weight (7)		≥ 2 Yrs Over-weight (7)			
Clinic	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Tmce-Van H	25	* *	* *	76	* *	* *	* *	30	* *	* *	51	* *
Tmcw-Ldrp	854	11.0 (88)	6.8 (51)	874	3.0 (3)	16.4 (16	4.6 (2)	51	* *	* *	96	* *
Tmcw-OB Cl	210	12.9 (100)	4.3 (17)	555	7.6 (10	7.7 (13	11.4 (58)	230	17.0 (77)	11.3 (35)	349	26.6 (142)
Tmcw-Pedia	98	* *	* *	605	12.4 (16	6.6 (13	15.0 (13	340	14.4 (34)	12.4 (57)	475	22.5 (129)
Tri-County	54	* *	* *	183	6.6 (76)	1.6 (13	14.8 (12	86	* *	* *	123	17.9 (108)
Vernon Co	167	6.6 (16)	7.8 (77)	640	6.1 (61)	3.4 (81)	12.5 (85)	280	15.7 (58)	13.9 (76)	447	11.4 (53)
Washington	253	13.8 (105)	6.7 (48)	899	11.6 (15	2.9 (57)	14.2 (11	375	21.1 (12	17.6 (11	621	7.7 (18)
Watertower	277	15.9 (110)	4.3 (17)	680	12.1 (15	10.3 (15	14.7 (12	230	17.8 (88)	21.7 (12	338	16.9 (97)
Wayne Co H	133	10.5 (81)	3.8 (9)	582	6.4 (67)	2.9 (57)	14.6 (12	316	17.1 (78)	19.0 (12	428	6.3 (10)
Webster Co	288	5.9 (10)	8.3 (89)	1,125	6.4 (67)	4.0 (98)	10.0 (34)	548	16.2 (63)	11.1 (32)	808	9.5 (37)
Wellston W	5	* *	* *	29	* *	* *	* *	15	* *	* *	22	* *

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 6A
2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 17
Run Date: 7 / 8 / 04

	Comparison of Growth and Anemia Indicators by Clinic (3,4)										
	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)	
	Low	High			Short Stature (6)	Under-weight (7)	Over-weight (7)	≥ 2 Yrs Over-weight (7)			
Clinic	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- <95th (Rank)	% >95th (Rank)	Number	% (Rank)
Wentzville	1	* *	* *	1	* *	* *	* *	0	* *	* *	0	* *
Worth-Tri	6	* *	* *	48	* *	* *	* *	28	* *	* *	38	* *
Wright Cou	168	9.5 (59)	4.8 (26)	744	6.0 (57)	2.7 (50)	10.9 (50)	359	16.4 (65)	13.9 (76)	537	15.1 (81)
Missouri	**39,423**	**9.1**	**6.9**	**131,869**	**6.9**	**4.6**	**12.0**	**56,346**	**16.4**	**13.3**	**86,726**	**16.2**
Nation (Prior Year)	**1,208,400**	**8.9**	**7.6**	**4,913,316**	**6.3**	**5.4**	**13.5**	**2,069,605**	**15.4**	**14.3**	**3,295,974**	**13.1**

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)					
	Breastfeeding (6)				
Clinic	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
(Delmar) P	602	37.4 (89)	59	* *	163	30.7 (27)
(Maplewood	160	51.9 (44)	29	* *	65	* *
00Family C	0	* *	0	* *	0	* *
00Miller C	38	* *	17	* *	18	* *
01SL Conne	135	17.0(110)	20	* *	57	* *
02Family C	157	51.0 (46)	41	* *	99	* *
Andrew Co	102	37.3 (91)	24	* *	49	* *
Atchison C	32	* *	29	* *	40	* *
Audrain Co	183	43.2 (76)	6	* *	46	* *
Barnes Hos	166	50.6 (47)	1	* *	0	* *
Barry Co H	385	53.2 (36)	73	* *	184	33.2 (20)
Barton Co	105	52.4 (42)	38	* *	83	* *
Bates Co H	132	48.5 (52)	36	* *	88	* *

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributers.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 7A
2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)					
	Breastfeeding (6)				
Clinic	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
BeltonCass	10	* *	9	* *	20	* *
Benton Co	104	43.3 (74)	39	* *	68	* *
Blue Sprin	367	63.8 (7)	74	* *	147	17.7 (55)
Bollinger	90	* *	4	* *	28	* *
BransTaney	248	68.1 (2)	111	29.7 (19)	219	26.0 (34)
Butler Co	430	44.0 (70)	41	* *	143	30.1 (28)
Callaway C	198	48.0 (53)	34	* *	98	* *
Camden Co	240	45.8 (62)	54	* *	113	18.6 (51)
Cape Girar	421	42.5 (79)	122	19.7 (26)	230	23.0 (43)
Carondelet	386	52.6 (40)	99	* *	187	39.6 (8)
Carroll Co	74	* *	16	* *	44	* *
Carter Co	48	* *	9	* *	22	* *
Cass Co He	321	58.9 (17)	70	* *	148	31.1 (24)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributers.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in
   the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 7A
2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)						
	Breastfeeding (6)					
Clinic	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Cedar Co H	115	53.9 (34)	11	* *	32	* *
Chariton C	36	* *	14	* *	26	* *
Christian	339	65.5 (4)	93	* *	202	13.4 (64)
Clark Co H	52	* *	22	* *	37	* *
Clay Chd-E	5	* *	0	* *	5	* *
Clay Chd-P	12	* *	3	* *	6	* *
Clay Co He	755	55.9 (27)	135	30.4 (17)	262	25.6 (35)
Clay Metro	2	* *	0	* *	2	* *
Clinton Co	154	53.2 (36)	44	* *	95	* *
Cole Co He	467	59.3 (16)	175	33.1 (13)	320	35.3 (18)
Columbia-B	554	61.2 (11)	144	45.1 (5)	284	36.3 (15)
Comp Healt	250	18.0(109)	9	* *	62	* *
Cooper Co	85	* *	5	* *	26	* *

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributers.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in
   the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)						
	Breastfeeding (6)					
Clinic	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Crawford C	207	46.4 (60)	23	* *	63	* *
Crescent C	413	60.0 (12)	129	38.0 (7)	250	41.6 (5)
Dade Co He	69	* *	20	* *	39	* *
Dallas Co	120	49.2 (51)	36	* *	78	* *
Daviess Co	72	* *	20	* *	40	* *
Dekalb Co-	35	* *	2	* *	10	* *
Dent Co He	118	43.2 (76)	39	* *	62	* *
Douglas Co	103	57.3 (22)	40	* *	67	* *
Dunklin Co	284	28.5(104)	62	* *	137	20.4 (48)
Elsber-Lnc	10	* *	0	* *	3	* *
Florence H	257	38.9 (87)	74	* *	158	23.4 (41)
Forest Par	380	31.6(101)	53	* *	135	31.1 (24)
Franklin C	462	43.9 (71)	105	36.2 (9)	241	25.3 (36)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributers.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 7A
2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)					
	Breastfeeding (6)				
Clinic	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
FtWood-Pul	269	61.3 (10)	98	* *	185	35.7 (17)
Gasconade	100	50.0 (48)	40	* *	74	* *
GascondOsg	40	* *	13	* *	29	* *
Grace Hill	129	30.2(102)	27	* *	63	* *
Grundy Co	87	* *	21	* *	40	* *
Hamilton/C	25	* *	6	* *	17	* *
Harrison C	78	* *	24	* *	45	* *
Hartville-	34	* *	6	* *	17	* *
Henry Co H	154	56.5 (25)	72	* *	118	19.5 (50)
HGP-Connec	47	* *	5	* *	20	* *
Hickory Co	63	* *	13	* *	34	* *
Holt Count	30	* *	8	* *	16	* *
Hosphill-T	371	33.7 (98)	198	36.4 (8)	492	35.8 (16)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributers.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in
 the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 7A
2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)					
	Breastfeeding (6)				
Clinic	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Howard Co	51	* *	11	* *	31	* *
Iron Co He	98	* *	30	* *	63	* *
Jfcac-Arno	457	46.6 (59)	56	* *	159	18.2 (52)
Jfcac-Unit	543	43.8 (73)	111	22.5 (25)	252	22.6 (44)
John C Mur	1,031	35.5 (94)	183	24.0 (23)	341	22.0 (45)
Johnson Co	394	68.8 (1)	138	28.3 (20)	273	24.5 (38)
Joplin Cit	1,064	55.7 (28)	207	28.0 (22)	500	21.6 (46)
Kansas Cit	0	* *	0	* *	0	* *
Kchd-Ruski	0	* *	0	* *	0	* *
Knox Co He	24	* *	13	* *	15	* *
La Clinica	220	57.3 (22)	21	* *	83	* *
Laclede Co	317	28.7(103)	83	* *	180	27.8 (31)
Lakewood-T	537	52.7 (39)	154	47.4 (4)	272	39.0 (10)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributers.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in
    the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)					
	Breastfeeding (6)				
Clinic	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Lawrence C	303	41.3 (81)	114	28.1 (21)	229	28.8 (29)
Lewis Co H	73	* *	22	* *	47	* *
Lincoln Co	232	40.9 (83)	62	* *	124	16.1 (57)
Linn Co He	100	45.0 (66)	40	* *	63	* *
Livingston	111	47.7 (55)	51	* *	84	* *
Macon Co H	103	52.4 (42)	32	* *	66	* *
Madison Co	135	36.3 (92)	46	* *	95	* *
Malden-Dnk	128	27.3(105)	62	* *	106	15.1 (60)
Maries Co	39	* *	6	* *	15	* *
Marion Co	249	33.7 (98)	85	* *	191	27.7 (32)
Mcdonald C	315	44.1 (69)	100	24.0 (23)	191	20.9 (47)
Mercer Co	36	* *	9	* *	24	* *
Midtown WI	93	* *	11	* *	33	* *

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributers.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in
  the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)						
	Breastfeeding (6)					
Clinic	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Miller co	185	44.9 (67)	59	* *	106	31.1 (24)
Mississipp	158	26.6(106)	70	* *	125	16.8 (56)
Moniteau C	69	* *	29	* *	41	* *
Monroe Co	64	* *	5	* *	22	* *
Montgomery	78	* *	19	* *	44	* *
Morgan Co	140	42.9 (78)	57	* *	103	40.8 (7)
NE Communi	174	41.4 (80)	64	* *	117	32.5 (22)
New Madrid	179	25.7(107)	71	* *	107	14.0 (63)
Newton Co	354	63.6 (9)	108	31.5 (15)	257	24.1 (40)
Nmhc-WIC-A	166	63.9 (6)	37	* *	95	* *
Nodaway Co	101	54.5 (31)	21	* *	39	* *
North Coun	115	46.1 (61)	22	* *	34	* *
Oregon Co	111	57.7 (21)	35	* *	77	* *

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributers.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)					
	Breastfeeding (6)				
Clinic	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Ozark Co H	60	* *	26	* *	59	* *
Pemiscot C	286	9.1(112)	31	* *	122	4.9 (66)
Perry Co H	127	43.3 (74)	42	* *	78	* *
Pettis Cou	375	54.4 (33)	90	* *	182	15.9 (58)
Phelps Co	301	56.1 (26)	57	* *	149	18.1 (53)
Pike Co He	138	34.8 (96)	39	* *	85	* *
Pine Lawn	534	40.4 (84)	115	31.3 (16)	210	28.6 (30)
Platte Chd	329	65.7 (3)	97	* *	202	44.1 (3)
Platte Co	43	* *	5	* *	18	* *
Polk Co He	212	63.7 (8)	20	* *	68	* *
Polo-Braym	17	* *	4	* *	15	* *
Prince Hal	154	39.6 (85)	32	* *	70	* *
Pulaski Co	141	47.5 (56)	38	* *	85	* *

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributers.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in
    the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2003 Pediatric Nutrition Surveillance [1]
Missouri
Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic [2]
Children Aged < 5 Years [3]

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic [4,5]					
	Breastfeeding [6]				
Clinic	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Putnam Co	32	* *	6	* *	17	* *
Ralls Co H	42	* *	12	* *	17	* *
Randolph C	209	51.2 (45)	34	* *	93	* *
Ray Co Hea	152	46.7 (58)	49	* *	80	* *
Reynolds C	55	* *	6	* *	28	* *
Richard Ca	31	* *	3	* *	21	* *
Ripley Co	115	53.0 (38)	37	* *	68	* *
Saline Co	172	58.7 (18)	36	* *	64	* *
Sam-Rod-Hi	46	* *	18	* *	28	* *
Sam-Rod-Le	60	* *	14	* *	25	* *
Sam-Rod-Od	44	* *	18	* *	38	* *
Samuel U R	594	59.9 (14)	220	53.6 (1)	417	53.0 (1)
Schuyler C	25	* *	15	* *	29	* *

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributers.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in
   the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 7A
2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)					
	Breastfeeding (6)				
Clinic	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Scotland C	28	* *	7	* *	12	* *
Scott Co H	439	39.4 (86)	146	17.1 (27)	277	18.1 (53)
Sears Cert	390	35.4 (95)	56	* *	150	32.7 (21)
Shannon Co	80	* *	28	* *	60	* *
Shelby Co	52	* *	18	* *	29	* *
SO Central	394	50.0 (48)	149	43.0 (6)	284	37.3 (13)
Soulard/Gr	131	41.2 (82)	27	* *	66	* *
South Coun	749	44.2 (68)	175	30.3 (18)	311	24.8 (37)
South Grac	177	49.7 (50)	35	* *	60	* *
South Poin	423	37.4 (89)	75	* *	206	37.9 (12)
Southside	21	* *	1	* *	7	* *
Spgfld-Gre	1,797	55.1 (30)	510	35.7 (10)	1,048	38.5 (11)
St Charles	590	59.7 (15)	48	* *	204	24.5 (38)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributers.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 7A
2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)						
	Breastfeeding (6)					
Clinic	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
St Clair C	65	* *	17	* *	35	* *
St Francoi	480	33.5(100)	32	* *	120	14.2 (62)
St Joseph-	627	37.5 (88)	50	* *	176	10.8 (65)
St Louis C	987	43.9 (71)	188	35.1 (11)	319	34.2 (19)
St Louis H	0	* *	0	* *	1	* *
St Lukes N	373	45.3 (64)	66	* *	171	41.5 (6)
St Stephen	4	* *	0	* *	0	* *
Starkloff-	202	36.1 (93)	29	* *	78	* *
StCHarvstr	37	* *	1	* *	10	* *
StChrlsVFW	91	* *	9	* *	21	* *
Ste Genevi	95	* *	30	* *	56	* *
Stoddard C	226	34.5 (97)	9	* *	66	* *
Stone Co H	171	52.6 (40)	48	* *	103	36.9 (14)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributers.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in
   the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)						
	Breastfeeding (6)					
Clinic	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Sullivan C	70	* *	24	* *	42	* *
Swope Pkwy	776	47.4 (57)	89	* *	251	15.1 (60)
Taney Coun	138	64.5 (5)	64	* *	122	39.3 (9)
Texas Co H	180	60.0 (12)	74	* *	141	31.9 (23)
Tmce-Grand	517	58.6 (19)	169	32.5 (14)	321	26.2 (33)
Tmce-Jacks	660	53.8 (35)	202	33.7 (12)	421	20.2 (49)
Tmce-Lee's	0	* *	0	* *	1	* *
Tmce-Oak G	72	* *	28	* *	47	* *
Tmce-Susqu	43	* *	17	* *	37	* *
Tmce-Van H	25	* *	10	* *	14	* *
Tmcw-Ldrp	871	48.0 (53)	10	* *	24	* *
Tmcw-OB Cl	207	55.6 (29)	41	* *	69	* *
Tmcw-Pedia	98	* *	23	* *	64	* *

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributers.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in
   the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 7A
2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)					
	Breastfeeding (6)				
Clinic	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Tri-County	54	* *	3	* *	18	* *
Vernon Co	164	56.7 (24)	36	* *	86	* *
Washington	253	17.0 (110)	60	* *	144	15.3 (59)
Watertower	277	24.5 (108)	47	* *	107	23.4 (41)
Wayne Co H	133	45.1 (65)	48	* *	94	* *
Webster Co	287	58.5 (20)	118	49.2 (2)	203	41.9 (4)
Wellston W	5	* *	0	* *	1	* *
Wentzville	1	* *	1	* *	0	* *
Worth-Tri	6	* *	0	* *	4	* *
Wright Cou	166	45.8 (62)	33	* *	74	* *
Unknown Clinics	367	54.5 (31)	105	48.6 (3)	198	49.0 (2)
Missouri	**38,757**	**47.7**	**9,108**	**32.4**	**19,376**	**28.6**

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributers.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 7A
2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)						
	Breastfeeding (6)					
Clinic	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Nation (Prior Year)	**969,228**	**52.5**	**527,765**	**20.8**	**555,990**	**12.3**

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributers.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in
the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 6A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 1
Run Date: 7 / 8 / 04

Comparison of Growth and Anemia Indicators by Clinic (3,4)

Clinic	Birthweight (5) Low Number	% <2500 g (Rank)	High % >4000 g (Rank)	Height and Weight Number	Short Stature (6) % <5th (Rank)	Under-weight (7) % <5th (Rank)	Over-weight (7) % >95th (Rank)	≥ 2 Yrs Over-weight (7) Number	% 85th- <95th (Rank)	% >95th (Rank)	Anemia Low Hb/Hct (8) Number	% (Rank)
(Delmar) P	1,743	11.7 (133)	4.2 (15)	4,507	7.5(13	10.9(17	10.0(40)	1,642	14.0(38)	13.2(97)	2,681	27.6(163)
(Maplewood	607	12.4 (144)	6.4 (44)	1,781	8.8(15	8.1(15	15.1(15	692	18.1(13	21.5(16	1,117	23.9(152)
00Family C	0	* *	* *	1	* *	* *	* *	1	* *	* *	1	* *
00Miller C	105	6.7 (23)	7.6 (76)	358	5.3(31)	1.7 (6)	11.5(79)	148	14.2(44)	18.2(16	239	13.0 (73)
01SL Conne	389	11.8 (134)	5.4 (30)	1,216	9.1(16	8.2(15	17.4(17	485	23.7(16	20.8(16	791	12.6 (70)
01St Louis	2	* *	* *	3	* *	* *	* *	1	* *	* *	1	* *
02Family C	487	13.3 (153)	4.7 (20)	1,626	8.2(14	8.9(15	11.9(89)	676	14.8(57)	11.5(53)	1,134	18.4 (124)
Andrew Co	262	6.1 (11)	7.6 (76)	1,022	4.4(13	5.4(13	11.2(68)	483	15.9(88)	12.8(87)	692	3.8 (2)
Atchison C	115	5.2 (5)	7.8 (83)	526	6.7(10	4.2(11	16.7(17	253	18.2(13	17.4(15	406	23.2 (147)
Audrain Co	485	7.8 (55)	6.4 (44)	1,546	9.8(16	2.1(17	14.4(14	615	16.6(10	13.2(97)	1,051	17.4 (116)
Barnes Hos	454	10.1 (109)	5.9 (39)	436	6.0(65)	15.1(17	3.0 (1)	1	* *	* *	3	* *

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 6A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low		High		Short Stature (6)	Under-weight (7)	Over-weight (7)		≥ 2 Yrs Over-weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- <95th (Rank)	% >95th (Rank)	Number	% (Rank)
Barry Co H	1,129	8.1 (64)	6.8 (51)	4,205	6.3 (79)	3.7 (87)	10.8 (59)	1,813	14.9 (59)	13.0 (92)	2,858	11.7 (59)
Barton Co	334	6.0 (9)	7.5 (73)	1,452	6.7 (10	2.3 (23)	13.6 (13	660	21.1 (16	15.6 (14	1,018	14.7 (92)
Bates Co H	405	7.4 (43)	5.9 (39)	1,554	6.6 (94)	5.6 (13	12.0 (93)	698	21.9 (16	16.3 (14	1,061	20.4 (135)
BeltonCass	53	* *	* *	417	6.5 (89)	3.1 (64)	12.2 (10	220	17.7 (12	11.4 (52)	324	27.2 (162)
Benton Co	319	8.5 (75)	7.8 (83)	1,280	6.9 (10	5.9 (14	9.6 (28)	616	17.2 (11	14.8 (12	889	13.7 (79)
Blue Sprin	1,001	8.9 (87)	7.5 (73)	3,546	5.9 (59)	4.4 (11	9.6 (28)	1,479	16.4 (10	10.3 (30)	2,347	12.2 (67)
Bollinger	287	8.4 (73)	5.6 (34)	1,111	5.8 (52)	3.5 (77)	10.3 (47)	488	15.8 (86)	12.7 (84)	781	10.6 (42)
BransTaney	789	8.0 (59)	9.0 (133)	3,159	5.5 (39)	2.7 (48)	11.1 (65)	1,469	16.7 (10	11.7 (59)	2,217	10.4 (38)
Butler Co	1,255	9.2 (94)	5.3 (29)	4,585	10.0 (17	1.8 (7)	14.4 (14	1,977	18.7 (14	16.3 (14	3,052	7.2 (20)
Callaway C	548	12.4 (144)	6.9 (54)	1,802	8.3 (14	3.2 (70)	11.4 (75)	724	17.1 (11	12.2 (69)	1,194	10.9 (47)
Camden Co	702	7.0 (34)	7.0 (57)	2,492	4.5 (16)	1.8 (7)	8.0 (10)	1,053	13.7 (34)	8.2 (9)	1,665	9.1 (27)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 6A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low		High		Short Stature (6)	Under-weight (7)	Over-weight (7)		≥ 2 Yrs Over-weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Cape Girar	1,267	8.5 (75)	6.6 (48)	3,765	8.2 (14	3.1 (64)	12.7 (11	1,338	17.5 (12	13.3 (10	2,364	21.2 (137)
Carondelet	1,085	10.1 (109)	7.4 (69)	3,539	6.8 (10	8.6 (15	11.9 (89)	1,444	14.3 (46)	13.4 (10	2,373	9.4 (31)
Carroll Co	230	5.2 (5)	7.0 (57)	922	4.4 (13)	2.9 (53)	12.0 (93)	434	17.3 (11	11.1 (47)	653	9.3 (30)
Carter Co	166	9.0 (90)	2.4 (1)	807	4.5 (16)	7.2 (14	7.3 (5)	411	10.7 (3)	6.6 (2)	591	16.8 (112)
Cass Co He	969	7.7 (52)	8.8 (120)	3,489	6.5 (89)	4.3 (11	10.8 (59)	1,517	13.4 (28)	9.2 (17)	2,274	19.2 (131)
Cedar Co H	336	6.8 (26)	10.4 (151)	1,449	5.5 (39)	2.9 (53)	13.9 (13	705	18.9 (14	17.0 (15	1,044	14.5 (88)
Chariton C	108	6.5 (16)	8.3 (106)	459	4.1 (10)	2.0 (13)	12.6 (11	213	17.4 (12	9.9 (24)	321	5.3 (4)
Christian	1,053	6.0 (9)	7.9 (91)	4,167	6.2 (75)	3.8 (92)	10.6 (56)	1,883	15.6 (80)	11.0 (45)	2,905	12.3 (69)
Clark Co H	168	11.9 (136)	9.5 (142)	591	6.1 (68)	2.9 (53)	10.3 (47)	257	17.9 (12	16.7 (15	392	5.6 (8)
Clay Chd-E	30	* *	* *	205	8.8 (15	2.9 (53)	14.1 (13	112	23.2 (16	12.5 (79)	153	15.7 (102)
Clay Chd-P	37	* *	* *	167	6.0 (65)	2.4 (29)	13.8 (13	78	* *	* *	121	14.9 (94)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 4
Run Date: 7 / 8 / 04

Comparison of Growth and Anemia Indicators by Clinic (3,4)

Clinic	Birthweight (5)		Height and Weight							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under-weight (7)	Over-weight (7)		≥ 2 Yrs Over-weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- <95th (Rank)	% >95th (Rank)	Number	% (Rank)
Clay Co He	2,187	8.7 (84)	8.5 (111)	6,420	8.5 (15	3.8 (92)	12.8 (11	2,456	17.1 (11	13.1 (94)	4,000	15.7 (102)
Clay Metro	14	* *	* *	103	3.9 (6)	0.0 (1)	18.4 (17	55	* *	* *	79	* *
Clinton Co	428	8.2 (66)	9.6 (144)	1,632	6.3 (79)	3.5 (77)	13.5 (12	731	16.8 (11	14.5 (12	1,154	7.1 (19)
Cochran He	16	* *	* *	73	* *	* *	* *	35	* *	* *	57	* *
Cole Co He	1,386	8.6 (81)	8.4 (109)	5,147	5.3 (31)	4.0 (10	10.3 (47)	2,288	15.6 (80)	12.6 (81)	3,523	6.8 (14)
Columbia-B	1,893	10.2 (113)	7.1 (61)	5,688	7.0 (11	3.7 (87)	11.2 (68)	2,122	16.5 (10	12.8 (87)	3,589	25.2 (157)
Comp Healt	767	10.6 (121)	4.3 (17)	2,156	6.3 (79)	8.9 (15	10.4 (52)	807	15.1 (64)	14.6 (12	1,407	21.5 (139)
Cooper Co	282	9.9 (104)	8.2 (104)	989	5.2 (28)	2.5 (35)	11.6 (80)	435	20.7 (16	13.6 (10	648	11.3 (55)
Crawford C	569	8.6 (81)	8.1 (103)	2,078	5.7 (46)	3.8 (92)	10.2 (44)	918	14.6 (53)	12.9 (90)	1,415	15.2 (98)
Crescent C	1,183	8.2 (66)	7.8 (83)	3,824	5.4 (36)	7.1 (14	10.0 (40)	1,522	12.7 (16)	8.7 (14)	2,569	38.0 (173)
Dade Co He	175	10.3 (115)	8.0 (96)	766	6.1 (68)	5.5 (13	10.3 (47)	370	16.2 (97)	13.2 (97)	533	11.8 (63)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)
	Low	High			Short Stature (6)	Under-weight (7)	Over-weight (7)	≥ 2 Yrs Over-weight (7)			
Clinic	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Dallas Co	327	9.2 (94)	8.6 (112)	1,161	7.7 (13	3.0 (59)	16.5 (16	505	15.4 (73)	14.9 (13	667	14.1 (83)
Daviess Co	184	7.6 (48)	7.6 (76)	771	9.2 (16	2.1 (17)	15.8 (16	383	13.1 (23)	14.6 (12	567	6.0 (10)
Dekalb Co-	116	9.5 (99)	6.0 (41)	499	6.4 (85)	3.6 (80)	11.2 (68)	249	15.3 (68)	10.0 (25)	349	16.3 (110)
Dent Co He	366	8.2 (66)	8.7 (114)	1,516	9.0 (16	3.2 (70)	9.3 (21)	708	15.0 (61)	7.6 (7)	1,073	9.2 (28)
Douglas Co	304	4.9 (4)	6.6 (48)	1,324	5.3 (31)	8.5 (15	10.0 (40)	662	15.3 (68)	10.4 (33)	926	14.0 (82)
Dunklin Co	866	9.0 (90)	4.0 (11)	3,087	7.0 (11	1.8 (7)	15.1 (15	1,345	18.5 (14	17.8 (15	2,145	11.1 (50)
Elsber-Lnc	92	* *	* *	296	10.1 (17	1.0 (3)	18.6 (17	127	15.0 (61)	13.4 (10	203	23.6 (150)
Florence H	793	14.0 (157)	4.2 (15)	2,616	8.9 (16	9.7 (16	12.9 (11	1,106	15.6 (80)	14.6 (12	1,661	14.1 (83)
Forest Par	844	10.5 (118)	5.7 (36)	1,401	5.2 (28)	12.0 (17	9.1 (19)	315	16.8 (11	16.5 (15	564	29.1 (166)
Franklin C	1,474	7.1 (36)	9.2 (138)	4,705	6.7 (10	3.6 (80)	11.1 (65)	1,787	15.2 (67)	11.1 (47)	3,081	10.0 (34)
FtWood-Pul	831	6.9 (32)	10.3 (149)	3,139	6.3 (79)	2.5 (35)	13.4 (12	1,475	11.3 (5)	9.2 (17)	2,276	31.4 (170)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 6A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 6
Run Date: 7 / 8 / 04

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)
	Low	High			Short Stature (6)	Under-weight (7)	Over-weight (7)	≥ 2 Yrs Over-weight (7)			
Clinic	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Gasconade	325	7.4 (43)	8.0 (96)	1,138	4.8 (20)	4.0 (10	9.6 (28)	484	15.3 (68)	14.0 (11	776	7.9 (22)
GascondOsg	125	4.8 (3)	11.2 (157)	442	6.8 (10	4.1 (10	11.8 (88)	202	20.8 (16	13.4 (10	304	10.9 (47)
Grace Hill	399	10.5 (118)	4.8 (21)	1,213	8.1 (14	8.2 (15	12.0 (93)	503	12.7 (16	13.9 (11	763	24.2 (153)
Grundy Co	273	6.6 (20)	5.5 (31)	1,056	6.1 (68	4.5 (12	12.1 (96)	489	18.2 (13	13.9 (11	737	7.9 (22)
Hamilton/C	69	* *	* *	310	6.1 (68)	1.3 (4)	15.5 (15	162	13.6 (31)	13.6 (10	230	4.3 (3)
Harrison C	228	9.2 (94)	10.5 (152)	900	4.0 (9)	2.4 (29)	12.8 (11	409	19.1 (15	13.9 (11	643	14.8 (93)
Hartville-	129	3.9 (1)	9.3 (139)	509	3.3 (2)	3.7 (87)	9.6 (28)	243	14.4 (47)	7.4 (6)	364	17.9 (119)
Henry Co H	517	8.3 (69)	8.9 (123)	1,864	5.9 (59)	3.6 (80)	11.7 (85)	780	15.0 (61)	10.3 (30)	1,265	17.1 (114)
HGP-Connec	207	11.6 (130)	3.4 (5)	595	11.8 (17	7.6 (14	13.1 (12	213	19.2 (15	16.4 (14	351	14.5 (88)
Hickory Co	156	9.0 (90)	7.7 (82)	542	6.6 (94)	4.1 (10	11.3 (74)	250	11.6 (9)	12.0 (64)	369	11.7 (59)
Holt Count	80	* *	* *	326	1.8 (1)	3.1 (64)	14.7 (14	156	17.9 (12	17.3 (15	237	14.3 (85)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 6A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)

Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low	High			Short Stature (6)	Under-weight (7)	Over-weight (7)	≥ 2 Yrs Over-weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- <95th (Rank)	% >95th (Rank)	Number	% (Rank)
Hosphill-T	1,682	11.6 (130)	6.0 (41)	8,452	7.6 (13	4.7 (12	12.7 (11	3,765	16.5 (10	11.5 (53)	6,388	23.8 (151)
Howard Co	149	12.8 (148)	8.7 (114)	514	4.7 (19)	2.9 (53)	7.6 (7)	216	10.6 (2)	6.5 (1)	353	10.5 (41)
Iron Co He	302	9.3 (97)	9.6 (144)	1,208	5.8 (52)	2.6 (45)	12.7 (11	571	17.5 (12	14.5 (12	860	6.4 (12)
Jeff-Frank	5	* *	* *	20	* *	* *	* *	11	* *	* *	13	* *
Jfcac-Arno	1,301	8.0 (59)	8.9 (123)	3,914	5.6 (43)	3.0 (59)	9.7 (33)	1,457	16.3 (10	13.6 (10	2,369	12.2 (67)
Jfcac-Unit	1,553	8.5 (75)	8.0 (96)	4,761	6.6 (94)	3.6 (80)	8.9 (15)	1,806	13.0 (20)	11.1 (47)	2,959	15.1 (97)
John C Mur	3,006	10.4 (117)	5.5 (31)	7,330	6.6 (94)	9.2 (16	8.0 (10)	2,468	12.6 (12)	9.6 (21)	4,316	22.6 (145)
Johnson Co	1,080	6.7 (23)	7.6 (76)	3,862	6.4 (85)	3.2 (70)	11.2 (68)	1,626	14.0 (38)	10.6 (37)	2,599	18.9 (127)
Joplin Cit	3,397	8.1 (64)	7.9 (91)	11,530	6.6 (94)	3.1 (64)	10.7 (58)	4,750	17.0 (11	11.9 (62)	7,521	18.1 (121)
Kansas Cit	444	8.8 (86)	8.6 (112)	1,744	7.5 (13	4.8 (12	11.4 (75)	721	14.1 (43)	11.8 (60)	1,266	22.7 (146)
Kchd-Clust	3	* *	* *	2	* *	* *	* *	0	* *	* *	0	* *

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 6A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)
	Low	High			Short Stature (6)	Under-weight (7)	Over-weight (7)	≥ 2 Yrs Over-weight (7)			
Clinic	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- <95th (Rank)	% >95th (Rank)	Number	% (Rank)
Kchd-Ruski	25	* *	* *	111	7.2(12	4.5(12	9.9(37)	59	* *	* *	83	* *
Knox Co He	80	* *	* *	454	4.4(13	2.6(45)	12.3(10	250	18.4(14	16.4(14	338	6.2 (11)
La Clinica	548	13.0 (151)	6.9 (54)	1,914	7.2(12	9.6(16	12.5(10	863	14.7(55)	15.5(13	1,370	26.8 (160)
Laclede Co	895	7.3 (40)	8.9(123)	2,921	7.5(13	4.1(10	11.0(63)	1,184	18.2(13	12.6(81)	1,851	10.7 (44)
Lakewood-T	1,609	6.8 (26)	8.3(106)	4,220	6.2(75)	5.8(13	11.2(68)	1,453	18.2(13	11.6(57)	2,439	18.9(127)
Lawrence C	915	7.3 (40)	7.9 (91)	3,615	5.3(31)	6.3(14	9.0(17)	1,683	11.3 (5)	8.3(11)	2,445	13.6 (77)
Lewis Co H	211	7.6 (48)	9.0(133)	926	4.8(20)	3.3(75)	13.2(12	477	15.5(76)	14.5(12	661	11.6 (58)
Lincoln Co	641	8.9 (87)	8.0 (96)	1,999	8.6(15	2.3(23)	14.2(14	774	16.5(10	13.6(10	1,308	14.5 (88)
Linn Co He	300	9.7(101)	5.0 (26)	1,066	6.9(10	0.9 (2)	16.6(16	462	19.5(15	19.9(16	729	13.7 (79)
Livingston	325	7.1 (36)	6.5 (47)	1,176	7.0(11	2.5(35)	14.4(14	499	14.0(38)	14.8(12	809	7.9 (22)
Macon Co H	314	9.6(100)	7.3 (65)	1,194	8.3(14	2.5(35)	16.2(16	528	22.0(16	18.8(16	817	10.2 (36)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 6A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 9
Run Date: 7 / 8 / 04

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)
	Low	High			Short Stature (6)	Under-weight (7)	Over-weight (7)	≥ 2 Yrs Over-weight (7)			
Clinic	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- <95th (Rank)	% >95th (Rank)	Number	% (Rank)
Madison Co	386	7.5 (47)	9.1 (135)	1,621	5.9 (59)	4.0 (10	16.2 (16	779	20.3 (15	20.8 (16	1,151	10.8 (45)
Malden-Dnk	413	11.9 (136)	4.4 (19)	1,661	7.3 (12	2.7 (48)	15.2 (15	801	19.5 (15	15.4 (13	1,176	11.7 (59)
Maries Co	122	7.4 (43)	10.7 (154)	506	3.6 (3)	4.2 (11	14.8 (15	238	16.4 (10	13.0 (92	360	11.7 (59)
Marion Co	769	10.8 (125)	7.3 (65)	2,702	8.5 (15	3.6 (80)	13.9 (13	1,115	18.1 (13	11.5 (53)	1,732	15.5 (100)
Mcdonald C	862	7.3 (40)	8.7 (114)	3,150	8.8 (15	2.4 (29)	19.3 (17	1,420	17.0 (11	15.5 (13	2,124	13.7 (79)
Mercer Co	90	* *	* *	336	3.9 (6)	3.0 (59)	14.0 (13	139	26.6 (16	15.1 (13	226	19.0 (130)
Midtown WI	291	12.4 (144)	4.8 (21)	911	9.0 (16	7.7 (14	9.7 (33)	377	15.9 (88)	11.1 (47)	610	29.0 (165)
Miller co	485	8.0 (59)	6.8 (51)	1,990	7.8 (14	2.2 (19)	13.1 (12	963	13.9 (37)	12.3 (74)	1,415	19.4 (132)
Mississipp	507	12.4 (144)	2.8 (3)	2,312	8.2 (14	2.5 (35)	13.3 (12	1,175	16.0 (92)	14.3 (12	1,780	16.1 (107)
Moniteau C	212	8.0 (59)	11.3 (158)	830	6.1 (68)	1.9 (12	12.4 (10	353	17.3 (11	12.2 (69)	559	13.4 (74)
Monroe Co	190	6.8 (26)	8.9 (123)	835	5.6 (43)	3.2 (70)	13.7 (13	435	19.1 (15	15.2 (13	599	11.2 (51)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 6A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 10
Run Date: 7 / 8 / 04

Comparison of Growth and Anemia Indicators by Clinic (3,4)										
	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)
	Low	High		Short Stature (6)	Under-weight (7)	Over-weight (7)	≥ 2 Yrs Over-weight (7)			
Clinic	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- <95th (Rank)	% >95th (Rank)	Number	% (Rank)
Montgomery	242	6.6 (20)	7.0 (57)	856	6.2 (75)	4.3 (11	11.6 (80)	375	10.9 (4)	10.1 (27)	591	14.4 (86)
Morgan Co	410	5.6 (7)	11.0 (156)	1,651	6.4 (85)	3.0 (59)	11.6 (80)	774	13.8 (36)	10.7 (39)	1,120	17.1 (114)
NE Communi	486	9.7 (101)	8.8 (120)	1,670	5.8 (52)	2.9 (53)	11.6 (80)	705	15.5 (76)	14.2 (11	1,098	16.6 (111)
New Madrid	536	10.1 (109)	3.4 (5)	2,222	8.0 (14	2.4 (29)	16.7 (17	1,050	15.7 (83)	15.3 (13	1,559	21.9 (143)
Newton Co	1,196	6.1 (11)	9.1 (135)	4,575	4.8 (20)	3.8 (92)	9.4 (25)	2,083	12.6 (12)	10.9 (43)	3,144	10.4 (38)
Nmhc-WIC-A	511	8.0 (59)	8.2 (104)	1,941	5.7 (46)	5.2 (13	7.8 (8)	875	13.1 (23)	10.4 (33)	1,341	18.9 (127)
Nodaway Co	294	7.1 (36)	10.9 (155)	1,052	5.3 (31)	3.8 (92)	9.8 (36)	485	13.6 (31)	10.1 (27)	709	17.5 (117)
North Coun	392	13.8 (156)	6.6 (48)	924	6.9 (10	9.3 (16	11.4 (75)	332	13.6 (31)	15.7 (14	538	26.0 (159)
Oregon Co	325	6.5 (16)	8.9 (123)	1,686	3.7 (4)	4.0 (10	12.3 (10	946	18.0 (13	12.2 (69)	1,295	10.0 (34)
Ozark Co H	200	8.5 (75)	9.5 (142)	807	5.8 (52)	3.6 (80)	10.8 (59)	375	18.1 (13	12.3 (74)	573	11.2 (51)
Pemiscot C	919	9.9 (104)	4.0 (11)	3,651	7.6 (13	2.4 (29)	15.6 (16	1,739	18.5 (14	15.3 (13	2,657	23.3 (149)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 11
Run Date: 7 / 8 / 04

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)
	Low	High			Short Stature (6)	Under-weight (7)	Over-weight (7)	≥ 2 Yrs Over-weight (7)			
Clinic	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- <95th (Rank)	% >95th (Rank)	Number	% (Rank)
People's H	1	* *	* *	24	* *	* *	* *	11	* *	* *	23	* *
Perry Co H	351	7.1 (36)	8.8 (120)	1,471	5.4 (36)	2.0 (13)	13.2 (12	670	17.9 (12	14.8 (12	1,065	11.3 (55)
Pettis Cou	1,146	7.9 (57)	7.9 (91)	4,489	5.5 (39)	3.0 (59)	11.1 (65)	2,012	14.7 (55)	12.0 (64)	3,108	10.8 (45)
Phelps Co	991	10.0 (107)	8.0 (96)	3,455	8.8 (15	4.3 (11	12.4 (10	1,447	13.4 (28)	12.1 (67)	2,367	12.0 (65)
Pike Co He	400	6.8 (26)	7.3 (65)	1,457	6.7 (10	4.7 (12	11.2 (68)	623	14.6 (53)	9.8 (23)	993	19.7 (133)
Pine Lawn	1,674	12.8 (148)	4.8 (21)	4,075	6.9 (10	10.6 (16	9.1 (19)	1,410	13.0 (20)	12.6 (81)	2,439	21.4 (138)
Platte Chd	965	7.8 (55)	10.3 (149)	3,338	6.4 (85)	3.8 (92)	13.5 (12	1,379	15.9 (88)	13.6 (10	2,191	21.5 (139)
Platte Co	142	12.0 (139)	4.9 (25)	463	5.8 (52)	4.3 (11	16.4 (16	193	21.8 (16	23.8 (16	310	21.0 (136)
Polk Co He	589	6.8 (26)	8.7 (114)	2,434	5.9 (59)	2.5 (35)	10.2 (44)	1,167	16.2 (97)	10.3 (30)	1,749	11.3 (55)
Polo-Braym	48	* *	* *	202	9.9 (17	2.0 (13)	11.4 (75)	84	* *	* *	148	5.4 (5)
Prince Hal	491	13.2 (152)	4.1 (13)	1,781	6.1 (68)	8.8 (15	11.0 (63)	794	13.7 (34)	10.1 (27)	1,282	32.1 (172)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 6A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)
	Low	High			Short Stature (6)	Under-weight (7)	Over-weight (7)	≥ 2 Yrs Over-weight (7)			
Clinic	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- <95th (Rank)	% >95th (Rank)	Number	% (Rank)
Pulaski Co	419	7.6 (48)	6.0 (41)	1,455	8.1(14	1.8 (7)	14.2(14	630	18.6(14	14.1(11	957	21.8 (142)
Putnam Co	120	10.0 (107)	5.8 (38)	537	4.3(12	5.6(13	10.6(56)	276	17.8(12	13.8(11	400	2.5 (1)
Ralls Co H	121	11.6 (130)	9.9 (147)	407	6.9(10	2.0(13	16.2(16	186	12.9(19	16.7(15	278	6.8 (14)
Randolph C	746	8.4 (73)	7.2 (63)	3,144	5.4(36	2.7(48	11.7(85)	1,477	18.9(14	12.7(84)	2,179	14.4 (86)
Ray Co Hea	392	11.0 (128)	7.4 (69)	1,418	8.5(15	3.7(87	9.4 (25)	618	14.2(44	7.0 (3)	937	15.5 (100)
Reynolds C	174	12.1 (142)	7.5 (73)	669	7.5(13	3.9(99	12.1(96)	297	18.5(14	13.5(10	443	8.4 (26)
Richard Ca	443	5.9 (8)	13.3 (159)	1,735	5.1(25	1.4 (5)	17.9(17	733	16.1(94	19.9(16	1,267	9.9 (33)
Ripley Co	421	10.2 (113)	2.9 (4)	1,747	7.7(13	5.4(13	8.9(15	807	12.6(12	7.3 (5)	1,210	11.2 (51)
Saline Co	565	6.5 (16)	7.4 (69)	1,944	6.9(10	3.7(87	12.2(10	793	14.0(38	13.1(94	1,286	15.7 (102)
Sam-Rod-Hi	153	3.9 (1)	10.5 (152)	568	5.5(39	2.3(23	15.0(15	248	19.8(15	16.1(14	365	11.2 (51)
Sam-Rod-Le	176	8.5 (75)	5.7 (36)	677	7.2(12	2.5(35	16.2(16	312	17.3(11	16.3(14	464	12.1 (66)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)

Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low	High			Short Stature (6)	Under-weight (7)	Over-weight (7)	≥ 2 Yrs Over-weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- <95th (Rank)	% >95th (Rank)	Number	% (Rank)
Sam-Rod-Od	117	12.0 (139)	6.8 (51)	439	8.9 (16	2.5 (35)	12.1 (96)	183	15.3 (68)	12.0 (64)	294	15.0 (95)
Samuel U R	1,588	6.2 (13)	8.9 (123)	6,436	3.9 (6)	6.1 (14	11.9 (89)	3,087	12.5 (11)	11.0 (45)	4,635	19.7 (133)
Schuyler C	88	* *	* *	384	9.1 (16	5.2 (13	11.7 (85)	179	15.1 (64)	12.3 (74)	281	6.8 (14)
Scotland C	82	* *	* *	288	5.6 (43)	2.4 (29)	7.3 (5)	118	21.2 (16	7.6 (7)	205	9.8 (32)
Scott Co H	1,262	10.1 (109)	4.3 (17)	4,831	7.9 (14	2.3 (23)	10.5 (54)	2,235	16.6 (10	12.3 (74)	3,373	17.0 (113)
Sears Cert	1,300	13.5 (155)	3.8 (10)	4,516	5.7 (46)	8.3 (15	9.3 (21)	2,015	14.0 (38)	10.0 (25)	3,161	29.1 (166)
Shannon Co	253	8.7 (84)	8.3 (106)	1,138	6.9 (10	2.5 (35)	14.0 (13	555	12.8 (18)	12.4 (78)	831	11.9 (64)
Shelby Co	169	10.7 (123)	4.1 (13)	780	5.8 (52)	2.3 (23)	13.3 (12	386	15.8 (86)	12.2 (69)	589	8.3 (25)
SO Central	1,058	6.9 (32)	9.4 (141)	4,246	3.7 (4)	4.6 (12	10.3 (47)	1,956	14.4 (47)	8.7 (14)	2,965	6.8 (14)
Soulard/Gr	416	10.8 (125)	7.2 (63)	1,308	10.8 (17	8.2 (15	17.9 (17	523	14.9 (59)	19.3 (16	882	16.2 (108)
South Coun	1,993	9.4 (98)	8.9 (123)	4,791	5.0 (23)	9.4 (16	8.8 (14	1,552	16.0 (92)	14.2 (11	2,710	15.9 (106)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 6A

2001 - 2003 Pediatric Nutrition Surveillance (1)

Missouri

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Page: 14

Run Date: 7 / 8 / 04

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)				
Clinic	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- <95th (Rank)	% >95th (Rank)	Number	% (Rank)
South Grac	342	10.8 (125)	5.6 (34)	585	10.3(17	11.3(17	15.2(15	115	18.3(14	30.4(16	225	24.9 (156)
South Poin	1,454	12.2 (143)	5.5 (31)	4,976	5.9(59	7.5(14	12.5(10	2,105	14.5(51	15.0(13	3,485	41.4 (174)
Southside	172	7.0 (34)	7.6 (76)	1,095	5.9(59	4.7(12	14.2(14	594	15.3(68	14.5(12	856	29.8 (168)
Spgfld-Gre	5,068	7.6 (48)	8.7 (114)	17,208	7.5(13	3.3(75	12.4(10	7,167	15.5(76	12.2(69	11,278	13.4 (74)
St Charles	2,158	8.3 (69)	7.6 (76)	5,698	6.3(79	3.5(77	9.9 (37)	2,058	16.4(10	11.8(60	3,262	7.6 (21)
St Clair C	218	13.3 (153)	7.3 (65)	1,035	8.7(15	3.2(70	12.1(96	539	19.3(15	12.1(67	783	13.4 (74)
St Francoi	1,395	8.3 (69)	8.9 (123)	4,755	6.1(68	4.0(10	9.4(25	1,907	15.9(88	11.6(57	3,129	14.6 (91)
St Joseph-	1,849	7.7 (52)	7.8 (83)	5,443	6.5(89	3.8(92	10.9(62	2,112	16.1(94	13.1(94	3,410	15.0 (95)
St Louis C	2,978	9.8 (103)	7.8 (83)	6,405	5.1(25	12.2(17	6.7 (4)	1,891	13.2(26	10.4(33	3,375	24.5 (155)
St Louis H	14	* *	* *	79	* *	* *	* *	49	* *	* *	66	* *
St Lukes N	992	10.3 (115)	7.4 (69)	2,644	7.6(13	5.0(12	14.1(13	917	15.4(73	10.7(39	1,552	16.2 (108)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 6A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 15
Run Date: 7 / 8 / 04

Comparison of Growth and Anemia Indicators by Clinic (3,4)

| Clinic | Birthweight (5) | | Height and Weight | | | | | | | Anemia Low Hb/Hct (8) | |
| | Low | High | | Short Stature (6) | Under-weight (7) | Over-weight (7) | | ≥ 2 Yrs Over-weight (7) | | | |
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- <95th (Rank)	% >95th (Rank)	Number	% (Rank)
St Stephen	54	* *	* *	156	9.0(16	11.5(17	15.4(15	64	* *	* *	96	* *
Starkloff-	553	11.9 (136)	3.4 (5)	1,517	6.3(79)	12.4(17	8.7(13)	561	11.4 (7)	8.4 (12)	852	10.6 (42)
StCHarvstr	112	6.3 (15)	3.6 (8)	364	6.9(10	2.2(19)	6.3 (3)	143	12.6(12)	7.0 (3)	182	5.5 (6)
StChrlsVFW	239	7.9 (57)	7.9 (91)	840	6.5(89)	3.1(64)	9.6(28)	348	14.4(47)	9.5(20)	488	5.5 (6)
Ste Genevi	287	6.6 (20)	9.1 (135)	1,081	7.7(13	2.5(35)	9.7(33)	502	17.9(12	10.6(37)	717	5.6 (8)
Stoddard C	705	10.6 (121)	4.8 (21)	2,572	7.3(12	4.1(10	12.4(10	1,096	17.8(12	14.1(11	1,760	27.6 (163)
Stone Co H	503	6.8 (26)	8.0 (96)	1,999	5.2(28)	4.1(10	7.9 (9)	891	13.0(20)	8.2 (9)	1,355	12.8 (71)
Sullivan C	200	11.5 (129)	7.0 (57)	766	5.1(25)	3.1(64)	16.7(17	327	18.3(14	16.8(15	484	10.3 (37)
Swope Pkwy	2,333	11.8 (134)	5.1 (28)	6,417	7.4(12	6.7(14	9.0(17)	2,348	13.1(23)	9.3(19)	4,159	22.3 (144)
Taney Coun	420	8.3 (69)	6.4 (44)	1,892	7.4(12	2.2(19)	10.4(52)	945	13.4(28)	10.7(39)	1,397	15.7 (102)
Texas Co H	577	9.0 (90)	8.7 (114)	2,329	4.5(16)	3.9(99)	9.9(37)	1,086	12.0(10)	10.9(43)	1,638	6.8 (14)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 6A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)

Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low		High		Short Stature (6)	Under-weight (7)	Over-weight (7)	≥ 2 Yrs Over-weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- <95th (Rank)	% >95th (Rank)	Number	% (Rank)
Tmce-Grand	1,431	8.6 (81)	7.8 (83)	5,132	6.7 (10	4.4 (11	12.4 (10	2,144	16.1 (94)	12.5 (79)	3,460	18.7 (126)
Tmce-Jacks	2,099	8.5 (75)	8.0 (96)	7,776	6.6 (94)	3.9 (99)	11.6 (80)	3,336	15.7 (83)	11.5 (53)	5,251	15.3 (99)
Tmce-Lee's	69	* *	* *	308	6.2 (75)	5.5 (13	12.7 (11	148	18.9 (14	12.8 (87)	215	27.0 (161)
Tmce-Oak G	202	9.9 (104)	8.9 (123)	791	7.0 (11	5.1 (13	13.8 (13	369	15.7 (83)	11.1 (47)	540	18.0 (120)
Tmce-Susqu	191	10.5 (118)	8.4 (109)	762	6.0 (65)	4.9 (12	10.5 (54)	361	13.3 (27)	8.9 (16)	520	17.5 (117)
Tmce-Van H	48	* *	* *	167	9.0 (16	1.8 (7)	15.6 (16	81	* *	* *	115	21.7 (141)
Tmcw-Ldrp	2,050	12.0 (139)	7.1 (61)	2,074	4.1 (10)	13.9 (17	4.0 (2)	139	11.5 (8)	10.8 (42)	226	24.3 (154)
Tmcw-OB Cl	441	12.9 (150)	5.0 (26)	1,109	7.3 (12	9.6 (16	9.3 (21)	448	14.5 (51)	8.5 (13)	685	31.5 (171)
Tmcw-Pedia	369	8.9 (87)	8.9 (123)	1,812	10.5 (17	6.3 (14	13.4 (12	929	14.4 (47)	12.7 (84)	1,354	25.9 (158)
Tri-Buchan	4	* *	* *	20	* *	* *	* *	11	* *	* *	16	* *
Tri-County	150	6.7 (23)	9.3 (139)	581	5.0 (23)	2.2 (19)	14.1 (13	272	16.2 (97)	12.9 (90)	399	18.3 (123)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)
		Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)			
Clinic	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Valley Par	188	17.6 (159)	9.6 (144)	509	5.7 (46)	9.6 (16	10.2 (44)	169	14.8 (57)	13.6 (10	316	13.6 (77)
Vernon Co	504	6.2 (13)	9.9 (147)	1,954	6.8 (10	2.6 (45)	13.4 (12	854	15.1 (64)	14.9 (13	1,363	10.9 (47)
Washington	778	10.7 (123)	7.8 (83)	2,784	10.2 (17	3.6 (80)	14.5 (14	1,212	20.6 (15	17.0 (15	1,927	10.4 (38)
Watertower	985	16.0 (158)	3.7 (9)	2,330	10.9 (17	10.1 (16	12.9 (11	809	17.3 (11	16.4 (14	1,265	23.2 (147)
Wayne Co H	407	7.4 (43)	2.7 (2)	1,798	6.6 (94)	2.3 (23)	14.8 (15	907	15.4 (73)	17.3 (15	1,288	18.2 (122)
Webster Co	818	6.5 (16)	7.8 (83)	3,360	5.8 (52)	4.6 (12	9.3 (21)	1,577	16.8 (11	10.5 (36)	2,405	9.2 (28)
Wellston W	12	* *	* *	66	* *	* *	* *	33	* *	* *	51	* *
Wentzville	57	* *	* *	185	6.5 (89)	2.7 (48)	11.9 (89)	73	* *	* *	108	6.5 (13)
West End W	77	* *	* *	309	7.1 (12	10.7 (17	8.1 (12)	146	9.6 (1)	9.6 (21)	230	30.9 (169)
Worth-Tri	38	* *	* *	194	5.7 (46)	8.8 (15	14.9 (15	97	* *	* *	152	18.4 (124)
Wright Cou	481	7.7 (52)	6.9 (54)	2,054	5.7 (46)	2.8 (52)	10.1 (43)	985	15.5 (76)	11.9 (62)	1,471	12.9 (72)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

2001 - 2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)										
	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)
	Low	High		Short Stature (6)	Under-weight (7)	Over-weight (7)	≥ 2 Yrs Over-weight (7)			
Clinic	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- <95th (Rank)	% >95th (Rank)	Number % (Rank)
Missouri	116,491	9.1	7.3	391,029	6.7	4.8	11.5	164,261	15.6	12.5	259,011 16.5

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 7A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)						
	Breastfeeding (6)					
Clinic	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
(Delmar) P	1,793	34.3(147)	692	29.6 (55)	824	28.9 (54)
(Maplewood	636	51.6 (63)	302	21.9(106)	349	13.2(131)
00Miller C	116	45.7(102)	74	* *	74	* *
01SL Conne	394	25.6(154)	213	9.9(139)	244	8.2(144)
01St Louis	1	* *	0	* *	0	* *
02Family C	514	49.6 (76)	301	39.2 (16)	341	36.1 (22)
Andrew Co	292	36.6(138)	181	24.9 (74)	199	26.1 (66)
Atchison C	134	33.6(148)	104	22.1(105)	115	23.5 (75)
Audrain Co	516	39.7(129)	253	22.5(102)	289	18.0(107)
Barnes Hos	407	52.1 (61)	32	* *	29	* *
Barry Co H	1,227	55.6 (43)	712	30.2 (54)	763	28.4 (56)
Barton Co	381	56.4 (36)	297	35.0 (31)	314	32.5 (37)
Bates Co H	445	47.9 (91)	273	19.4(116)	324	16.7(115)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributers.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in
    the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 7A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)					
	Breastfeeding (6)				
Clinic	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
BeltonCass	77	* *	78	* *	94	* *
Benton Co	347	51.0 (64)	224	37.9 (20)	254	38.2 (18)
Blue Sprin	1,144	65.6 (5)	565	24.4 (79)	648	16.4(118)
Bollinger	287	40.8(122)	145	22.8 (98)	180	16.7(115)
BransTaney	891	65.5 (6)	607	36.4 (23)	689	32.2 (40)
Butler Co	1,340	43.7(113)	680	26.9 (66)	800	26.4 (65)
Callaway C	606	43.1(116)	323	18.6(120)	363	16.3(119)
Camden Co	807	54.3 (51)	430	23.0 (92)	484	13.0(133)
Cape Girar	1,429	41.5(120)	751	24.4 (79)	843	23.6 (74)
Carondelet	1,145	48.3 (87)	618	35.6 (26)	705	33.8 (29)
Carroll Co	246	47.2 (96)	149	14.1(133)	182	9.9(140)
Carter Co	176	55.7 (42)	109	29.4 (57)	136	16.9(112)
Cass Co He	1,087	54.4 (49)	601	37.4 (21)	689	34.3 (27)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributers.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in
the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)					
	Breastfeeding (6)				
Clinic	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Cedar Co H	351	53.3 (57)	181	17.7(124)	224	11.2(138)
Chariton C	124	58.1 (24)	88	* *	94	* *
Christian	1,196	66.1 (3)	711	26.7 (68)	817	20.3 (95)
Clark Co H	180	55.6 (43)	115	34.8 (32)	122	36.9 (21)
Clay Chd-E	40	* *	27	* *	35	* *
Clay Chd-P	47	* *	31	* *	36	* *
Clay Co He	2,413	56.6 (34)	1,071	22.6 (99)	1,194	16.8(113)
Clay Metro	12	* *	13	* *	21	* *
Clinton Co	489	52.4 (58)	304	27.6 (63)	343	22.4 (79)
Cochran He	10	* *	6	* *	6	* *
Cole Co He	1,538	54.3 (51)	964	38.9 (17)	1,111	38.7 (17)
Columbia-B	2,029	62.4 (12)	1,020	33.0 (40)	1,133	27.9 (57)
Comp Healt	786	21.6(160)	325	15.4(132)	379	13.2(131)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributers.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in
   the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
 *   Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 7A
2001 - 2003 Pediatric Nutrition Surveillance [1]
Missouri
Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic [2]
Children Aged < 5 Years [3]

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic [4,5]						
	Breastfeeding [6]					
Clinic	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Cooper Co	276	48.9 (83)	123	8.1(140)	151	4.6(149)
Crawford C	617	44.7(107)	308	16.2(127)	357	9.5(143)
Crescent C	1,385	57.6 (27)	733	41.2 (11)	832	40.3 (12)
Dade Co He	191	57.6 (27)	125	38.4 (19)	153	39.9 (14)
Dallas Co	368	48.9 (83)	221	25.8 (70)	256	25.0 (70)
Daviess Co	211	51.7 (62)	122	24.6 (76)	142	17.6(110)
Dekalb Co-	119	65.5 (6)	55	* *	70	* *
Dent Co He	414	41.5(120)	262	23.7 (86)	303	21.5 (84)
Douglas Co	334	50.0 (70)	214	17.8(123)	244	20.5 (94)
Dunklin Co	929	30.7(151)	542	17.9(122)	619	15.5(123)
Elsber-Lnc	75	* *	43	* *	53	* *
Florence H	823	36.7(137)	458	28.2 (60)	548	29.2 (53)
Forest Par	853	35.2(144)	242	30.6 (52)	254	30.3 (49)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributers.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in
   the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 7A
2001 - 2003 Pediatric Nutrition Surveillance [1]
Missouri
Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic [2]
Children Aged < 5 Years [3]

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic [4,5]						
	Breastfeeding [6]					
Clinic	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Franklin C	1,593	45.6(104)	927	24.4 (79)	920	22.0 (82)
FtWood-Pul	1,003	59.9 (19)	614	34.2 (33)	694	29.5 (52)
Gasconade	344	37.8(133)	228	25.0 (73)	254	24.8 (71)
GascondOsg	138	50.0 (70)	79	* *	94	* *
Grace Hill	424	22.4(159)	207	20.8(109)	242	21.1 (87)
Grundy Co	286	50.3 (68)	157	21.0(108)	182	13.7(128)
Hamilton/C	77	* *	50	* *	66	* *
Harrison C	246	49.6 (76)	154	36.4 (23)	193	33.2 (34)
Hartville-	135	48.1 (90)	89	* *	102	30.4 (47)
Henry Co H	562	54.3 (51)	360	22.5(102)	433	19.9 (97)
HGP-Connec	204	28.4(153)	112	18.8(118)	129	16.3(119)
Hickory Co	170	44.1(109)	92	* *	109	31.2 (43)
Holt Count	91	* *	58	* *	61	* *

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributers.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in
   the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
 *   Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 7A
2001 - 2003 Pediatric Nutrition Surveillance [1]
Missouri
Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic [2]
Children Aged < 5 Years [3]

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic [4,5]					
	Breastfeeding [6]				
Clinic	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Hosphill-T	2,151	36.2(142)	1,695	30.7 (51)	1,924	30.1 (50)
Howard Co	168	50.6 (66)	96	* *	110	34.5 (26)
Iron Co He	319	41.7(119)	204	12.7(136)	251	8.0(146)
Jeff-Frank	5	* *	2	* *	1	* *
Jfcac-Arno	1,419	48.2 (88)	626	24.0 (85)	711	20.0 (96)
Jfcac-Unit	1,778	43.2(115)	809	24.4 (79)	915	22.2 (81)
John C Mur	3,156	36.4(140)	1,215	22.9 (95)	1,409	18.9(101)
Johnson Co	1,234	67.7 (2)	759	28.1 (61)	884	18.4(104)
Joplin Cit	3,761	60.0 (18)	1,971	27.4 (64)	2,254	22.4 (79)
Kansas Cit	397	54.7 (48)	276	33.3 (38)	332	32.8 (35)
Kchd-Clust	1	* *	0	* *	0	* *
Kchd-Ruski	19	* *	14	* *	15	* *
Knox Co He	94	* *	62	* *	69	* *

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributers.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in
   the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)						
	Breastfeeding (6)					
Clinic	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
La Clinica	624	59.5 (20)	330	49.1 (8)	367	49.9 (5)
Laclede Co	1,005	30.0(152)	551	23.0 (92)	619	23.3 (76)
Lakewood-T	1,696	56.1 (38)	820	34.0 (35)	906	25.6 (69)
Lawrence C	1,003	46.9 (99)	676	30.9 (49)	695	29.6 (51)
Lewis Co H	240	50.0 (70)	133	16.5(125)	167	10.8(139)
Lincoln Co	746	47.9 (91)	351	20.5(111)	404	12.6(134)
Linn Co He	319	43.9(111)	206	22.3(104)	231	20.8 (92)
Livingston	352	49.1 (81)	228	35.1 (29)	259	32.4 (38)
Macon Co H	326	58.9 (21)	213	39.9 (14)	246	40.2 (13)
Madison Co	414	39.4(130)	265	30.9 (49)	313	30.4 (47)
Malden-Dnk	446	24.2(156)	305	13.8(135)	353	13.3(130)
Maries Co	132	43.9(111)	82	* *	98	* *
Marion Co	858	36.2(142)	609	27.3 (65)	639	26.1 (66)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributers.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in
    the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2001 - 2003 Pediatric Nutrition Surveillance [1]
Missouri
Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic [2]
Children Aged < 5 Years [3]

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic [4,5]						
	Breastfeeding [6]					
Clinic	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Mcdonald C	985	48.6 (86)	651	22.6 (99)	690	20.9 (91)
Mercer Co	102	49.0 (82)	74	* *	79	* *
Midtown WI	319	42.3(117)	127	32.3 (43)	162	34.6 (25)
Miller co	526	45.8(101)	322	32.0 (44)	377	33.7 (32)
Mississipp	547	23.8(157)	415	16.4(126)	479	16.5(117)
Moniteau C	247	60.3 (17)	157	24.2 (84)	178	18.5(103)
Monroe Co	205	56.6 (34)	99	* *	119	39.5 (15)
Montgomery	259	36.3(141)	153	21.6(107)	179	21.2 (86)
Morgan Co	462	44.2(108)	288	32.6 (41)	324	33.6 (33)
NE Communi	554	45.1(106)	324	30.6 (52)	368	28.5 (55)
New Madrid	570	23.2(158)	414	15.7(129)	429	14.7(125)
Newton Co	1,334	62.6 (11)	813	24.7 (75)	923	19.6 (99)
Nmhc-WIC-A	565	65.8 (4)	323	36.8 (22)	352	21.3 (85)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributers.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in
   the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 7A
2001 - 2003 Pediatric Nutrition Surveillance [1]
Missouri
Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic [2]
Children Aged < 5 Years [3]

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic [4,5]						
	Breastfeeding [6]					
Clinic	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Nodaway Co	321	55.5 (45)	176	28.4 (59)	193	19.2(100)
North Coun	399	44.1(109)	144	34.0 (35)	166	30.7 (45)
Oregon Co	360	58.3 (23)	252	23.0 (92)	300	14.0(126)
Ozark Co H	227	56.8 (33)	164	41.5 (10)	181	39.2 (16)
Pemiscot C	977	11.5(162)	539	7.4(141)	653	6.3(148)
People's H	3	* *	2	* *	4	* *
Perry Co H	390	40.0(128)	270	11.9(137)	329	7.9(147)
Pettis Cou	1,252	55.8 (40)	757	20.5(111)	854	13.8(127)
Phelps Co	1,060	55.8 (40)	589	22.9 (95)	670	12.2(136)
Pike Co He	433	40.6(125)	265	15.5(131)	301	9.6(142)
Pine Lawn	1,738	35.0(145)	711	23.5 (88)	809	22.5 (78)
Platte Chd	1,106	63.2 (10)	646	41.2 (11)	722	35.9 (23)
Platte Co	158	64.6 (8)	77	* *	91	* *

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributers.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in
   the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 7A
2001 - 2003 Pediatric Nutrition Surveillance [1]
Missouri
Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic [2]
Children Aged < 5 Years [3]

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic [4,5]						
	Breastfeeding [6]					
Clinic	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Polk Co He	638	63.8 (9)	332	23.2 (90)	426	18.8(102)
Polo-Braym	58	* *	41	* *	44	* *
Prince Hal	510	40.4(126)	265	26.8 (67)	313	21.1 (87)
Pulaski Co	469	47.8 (93)	268	31.0 (48)	308	27.6 (60)
Putnam Co	131	53.4 (56)	77	* *	100	27.0 (62)
Ralls Co H	129	40.3(127)	66	* *	79	* *
Randolph C	731	54.4 (49)	493	19.9(114)	538	12.3(135)
Ray Co Hea	451	40.8(122)	248	24.6 (76)	271	26.9 (63)
Reynolds C	176	52.3 (59)	88	* *	112	24.1 (73)
Richard Ca	420	74.8 (1)	326	51.8 (2)	387	52.2 (1)
Ripley Co	467	50.1 (69)	292	10.6(138)	335	9.9(140)
Saline Co	596	53.5 (55)	323	23.2 (90)	372	19.9 (97)
Sam-Rod-Hi	169	40.8(122)	104	27.9 (62)	116	26.7 (64)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributers.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in
   the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 7A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)					
	Breastfeeding (6)				
Clinic	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Sam-Rod-Le	201	46.8(100)	126	32.5 (42)	141	32.6 (36)
Sam-Rod-Od	146	45.2(105)	93	* *	96	* *
Samuel U R	1,817	61.9 (14)	1,174	54.2 (1)	1,342	51.3 (2)
Schuyler C	100	50.0 (70)	71	* *	86	* *
Scotland C	92	* *	52	* *	62	* *
Scott Co H	1,382	38.9(132)	853	18.4(121)	991	16.8(113)
Sears Cert	1,371	36.5(139)	665	33.7 (37)	815	30.6 (46)
Shannon Co	268	47.0 (97)	185	20.0(113)	231	17.3(111)
Shelby Co	182	47.8 (93)	125	28.8 (58)	142	27.5 (61)
SO Central	1,215	50.9 (65)	757	35.8 (25)	875	32.3 (39)
Soulard/Gr	444	45.7(102)	227	50.2 (4)	273	48.7 (6)
South Coun	2,235	49.5 (78)	916	25.4 (72)	976	20.8 (92)
South Grac	369	49.3 (80)	127	50.4 (3)	121	50.4 (3)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributers.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in
    the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
 *   Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 7A
2001 - 2003 Pediatric Nutrition Surveillance [1]
Missouri
Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic [2]
Children Aged < 5 Years [3]

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic [4,5]						
	Breastfeeding [6]					
Clinic	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
South Poin	1,490	37.0(136)	792	35.1 (29)	972	33.8 (29)
Southside	134	41.8(118)	68	* *	145	37.9 (19)
Spgfld-Gre	5,696	57.3 (29)	3,163	35.5 (27)	3,622	31.2 (43)
St Charles	2,214	57.0 (32)	941	26.1 (69)	1,041	21.6 (83)
St Clair C	246	56.1 (38)	161	16.1(128)	194	11.3(137)
St Francoi	1,477	32.1(149)	647	19.5(115)	766	18.0(107)
St Joseph-	1,967	37.2(135)	782	15.6(130)	939	15.8(122)
St Louis C	3,169	47.0 (97)	1,127	31.7 (45)	1,225	27.8 (59)
St Louis H	11	* *	8	* *	12	* *
St Lukes N	1,068	48.2 (88)	494	41.1 (13)	589	40.7 (11)
St Stephen	49	* *	19	* *	26	* *
Starkloff-	574	39.0(131)	238	23.5 (88)	280	18.2(105)
StCHarvstr	124	62.1 (13)	56	* *	65	* *

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributers.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

CDC

Table 7A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)						
	Breastfeeding (6)					
Clinic	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
StChrlsVFW	295	54.9 (47)	127	24.4 (79)	143	21.0 (90)
Ste Genevi	312	37.8(133)	179	24.6 (76)	203	24.6 (72)
Stoddard C	750	32.0(150)	361	18.8(118)	451	18.0(107)
Stone Co H	559	50.6 (66)	348	34.2 (33)	396	33.8 (29)
Sullivan C	261	34.9(146)	175	22.9 (95)	175	21.1 (87)
Swope Pkwy	2,422	47.4 (95)	1,042	20.8(109)	1,198	13.4(129)
Taney Coun	478	61.1 (15)	344	38.7 (18)	387	33.9 (28)
Texas Co H	634	57.1 (31)	429	33.1 (39)	499	31.7 (42)
Tmce-Grand	1,656	57.9 (25)	999	31.5 (46)	1,106	27.9 (57)
Tmce-Jacks	2,347	55.0 (46)	1,371	23.6 (87)	1,530	15.9(121)
Tmce-Lee's	63	* *	42	* *	51	* *
Tmce-Oak G	234	57.3 (29)	132	29.5 (56)	146	26.0 (68)
Tmce-Susqu	199	54.3 (51)	124	25.8 (70)	149	18.1(106)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributers.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in
   the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2001 - 2003 Pediatric Nutrition Surveillance [1]
Missouri
Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic [2]
Children Aged < 5 Years [3]

Page: 14
Run Date: 7 / 8 / 04

Clinic	| Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic [4,5]					
	| Breastfeeding [6]					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Tmce-Van H	54	* *	29	* *	27	* *
Tmcw-Ldrp	1,692	49.9 (74)	293	50.2 (4)	285	50.2 (4)
Tmcw-OB Cl	492	58.7 (22)	204	50.0 (7)	209	46.9 (8)
Tmcw-Pedia	414	43.7(113)	275	35.3 (28)	317	34.7 (24)
Tri-Buchan	2	* *	2	* *	4	* *
Tri-County	161	49.7 (75)	71	* *	96	* *
Valley Par	155	52.3 (59)	79	* *	102	37.3 (20)
Vernon Co	553	60.6 (16)	326	39.3 (15)	393	41.0 (9)
Washington	828	18.8(161)	459	13.9(134)	531	15.4(124)
Watertower	1,000	24.5(155)	393	22.6 (99)	444	23.2 (77)
Wayne Co H	425	48.9 (83)	321	31.5 (46)	350	32.0 (41)
Webster Co	918	57.7 (26)	615	43.4 (9)	723	40.9 (10)
Wellston W	15	* *	6	* *	8	* *

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributers.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Missouri
Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)					
	Breastfeeding (6)				
Clinic	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Wentzville	41	* *	21	* *	23	* *
West End W	56	* *	29	* *	46	* *
Worth-Tri	40	* *	26	* *	36	* *
Wright Cou	524	49.4 (79)	287	19.2(117)	340	8.2(144)
Unknown Clinics	643	56.3 (37)	229	50.2 (4)	236	47.5 (7)
Missouri	**126,492**	**48.4**	**67,523**	**28.6**	**77,044**	**25.2**

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributers.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in
   the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.